arXiv:1208.2840v1 [math.DS] 14 Aug 2012

CONVERSE KAM THEORY REVISITED

Lin Wang

Abstract. For an integrable Hamiltonian with d ($d \geq 2$) degrees of freedom, we show the conditions on perturbations, for which invariant tori can be destructed.

Contents

1 Introduction 2

2 Destruction of a Lagrangian torus 4
- 2.1 Case with 2 degrees of freedom 4
  - 2.1.1 Preliminaries of exact area-preserving twist map 4
    - 2.1.1.1 Minimal configuration 4
    - 2.1.1.2 Peierls's barrier 5
  - 2.1.2 C^∞ case 6
    - 2.1.2.1 Construction of the generating functions 6
    - 2.1.2.2 Estimate of lower bound of $P_{0^+}^{h_n}$ 8
    - 2.1.2.3 The approximation from $P_{0^+}^{h_n}$ to $P_{\omega}^{h_n}$ 9
    - 2.1.2.4 Proof of Theorem 2.1 17
  - 2.1.3 C^ω case 18
    - 2.1.3.1 Construction of the generating functions 18
    - 2.1.3.2 Proof of Theorem 2.11 20
- 2.2 Case with $d \geq 3$ degrees of freedom 22
  - 2.2.1 Preliminaries 22
  - 2.2.2 C^∞ case 23
    - 2.2.2.1 Destruction of Lagrangian torus with a special rotation vector 24
    - 2.2.2.2 Destruction of Lagrangian torus with an arbitrary rotation vector 31
  - 2.2.3 C^ω case 34
    - 2.2.3.1 Construction of H_n 34
    - 2.2.3.2 Melnikov function 35
    - 2.2.3.3 An approximation lemma 36
    - 2.2.3.4 Proof of Theorem 2.22 44

3 Destruction of all Lagrangian tori 47
- 3.1 A toy model 47
- 3.2 C^∞ case 49
- 3.3 An approximation lemma 50
- 3.4 C^ω case 52

References 54

Date: August 15, 2012

1. Introduction

By the Kolmogorov, Arnold and Moser (KAM) theory, we know that under certain non-degeneracy, most (full Lebesgue measure) invariant tori of an integrable Hamiltonian system are persisted under small perturbations. As the sizes of the perturbations increase, those persisted invariant tori are destructed progressively. The problems of determining the critical boundary between the persistence and destruction of invariant tori motivates so called converse KAM theory.

Roughly speaking, the converse KAM theory consists of two parts. The first part, with more physical flavor, is concerned about destruction of invariant tori under the perturbations with positive lower bound, which is started by the analysis of invariant circles of the generalized standard maps (see e.g. [Ci],[H2] and [Ma2]). Besides, numerical results go further than theoretical ones. More sharp boundaries are obtained by numerical methods (see e.g. [Gr] and [MP]). The second part is concerned about destruction of invariant tori under arbitrarily small perturbations in certain topology and it seems that numerical method is not as efficient as the first part. In this paper, we are devoted to develop the second part.

In 1962, Moser proved that the invariant circles with Diophantine rotation numbers of an integrable twist map is persisted under arbitrarily small perturbations in the C^{333} topology ([Mo1]). By the efforts of Moser, Rüssman, Herman and Pöschel ([H2, H3],[Mo2, Mo3],[P] and [R1, R2]), for Hamiltonian systems with d-degrees of freedom, it is obtained that certain invariant tori are persisted under arbitrarily small perturbations in the $C^{2d+\delta}$ topology, where δ is a small positive constant. Especially, Herman proved in [H3] that for twist maps on annulus, certain invariant circles can be persisted under arbitrarily small perturbations in the C^3 topology.

In contrast with the results on persistence of invariant tori, for exact area-preserving twist maps on annulus, it is proved by Herman in [H2] that invariant circles with given rotation numbers can be destructed by $C^{3-\delta}$ arbitrarily small C^∞ perturbations. For certain rotation numbers, it is obtained by Mather (resp. Forni) in [Ma5] (resp. [F]) that the invariant circles with those rotation numbers can be destroyed by small perturbations in finer topology respectively. More precisely, Mather considers Liouville rotation numbers and the topology of the perturbation induced by C^∞ metric. Forni is concerned about more special rotation numbers which can be approximated by rational ones exponentially and the topology of the perturbation induced by the supremum norm of C^ω (real-analytic) function. Bessi extended the result to the systems with multi-degrees of freedom. He found that the invariant Lagrangian torus with certain rotation vector can be destructed by an arbitrarily small C^ω perturbation for certain positive definite systems with multi-degrees of freedom in [Be], where Lagrangian torus is a natural analogy to invariant circle in multi-degrees of freedom (see Definition 3.1 below).

On the other hand, it is also proved by Herman in [H5] that all Lagrangian tori of an integrable symplectic twist map with $d \geq 1$ degrees of freedom can be destructed by $C^{d+2-\delta}$ arbitrarily small C^∞ perturbations of the generating function. Equivalently ([Go, Mo4]), it shows that all Lagrangian tori of an integrable Hamiltonian system with $d \geq 2$ degrees of freedom can be destructed by C^∞ perturbations which are arbitrarily small in the $C^{d+1-\delta}$ topology. Roughly speaking, there is a balance among the arithmetic property of the rotation vector, the regularity of the perturbation and its topology.

Comparing the results on both sides, it is natural to ask the following questions:

- for every given rotation vector ω, if the Lagrangian torus with ω can be destructed by an arbitrarily small C^∞ perturbation in the C^r topology, then what is the maximum of r?
- for every given rotation vector ω, if the Lagrangian torus with ω can be destructed by an arbitrarily small analytic perturbation in the C^r topology, then what is the maximum of r?
- if all of Lagrangian tori can be destructed by an arbitrarily small real-analytic perturbation in the C^r topology, then what is the maximum of r?

Based on [CW] and [W1, W2, W3, W4], we have the following theorems.

Theorem 1.1 *Given an integrable Hamiltonian* H_0 *with* d $(d \geq 2)$ *degrees of freedom and a rotation vector* ω*, there exists a sequence of* C^∞ *Hamiltonians* $\{H_n\}_{n\in\mathbb{N}}$ *such that* $H_n \to H_0$ *in* $C^{2d-\delta}$ *topology and the Hamiltonian flow generated by* H_n *does not admit the Lagrangian torus with the rotation vector* ω*.*

Theorem 1.2 *Given an integrable Hamiltonian* H_0 *with* d $(d \geq 2)$ *degrees of freedom and a rotation vector* ω*, there exists a sequence of* C^ω *Hamiltonians* $\{H_n\}_{n\in\mathbb{N}}$ *such that* $H_n \to H_0$ *in* $C^{d+1-\delta}$ *topology and the Hamiltonian flow generated by* H_n *does not admit the Lagrangian torus with the rotation vector* ω*.*

Theorem 1.3 *For an integrable Hamiltonian* H_0 *with* d $(d \geq 2)$ *degrees of freedom, all Lagrangian tori can be destructed by analytic perturbations which are arbitrarily small in the* $C^{d-\delta}$ *topology.*

For positive definite Hamiltonian systems with $d \geq 2$ degrees of freedom, together with Herman's result in [H5], we have the following Table 1.

r	C^∞	C^ω
Single Destruction	$2d-\delta$	$d+1-\delta$
Total Destruction	$d+1-\delta$	$d-\delta$

Table 1: Values of r for destruction of Lagrangian torus (tori) in the C^r topology

Unfortunately, except for the destruction of a Lagrangian torus by the C^∞ perturbations, we still don't know whether the other results are optimal. Some further developments of KAM theory are needed to verify the optimality.

This paper is outlined as follows. In Section 2, we consider the destruction of a Lagrangian torus with given rotation vector (Theorem 1.1 and Theorem 1.2). Based on the difference of topology of phase spaces between $d = 2$ and $d \geq 3$, we divide the arguments into two cases. In terms of the correspondence between exact area-preserving twist maps and Hamiltonian systems with 2 degrees of freedom, the problem on destruction of the Lagrangian torus for the Hamiltonian system is transformed into the one on destruction of the invariant circle for the twist map. Using variational method developed by Mather, a new proof of Herman's result ([H2]) is provided. Moreover, from Jackson's approximation, the result on C^ω perturbation is obtained. For the case with $d \geq 3$ degrees of freedom, the minimality of the orbits on Lagrangian torus plays a crucial role in destruction of the torus. Combining with Melnikov method, the destruction of Lagrangian torus under C^ω perturbations is achieved. In Section 3, we are concerned about the destruction of all Lagrangian tori (Theorem 1.3). Similar to Section 2, according to the correspondence between the Hamiltonian system and the symplectic twist map, we focus on destruction of Lagrangian tori for symplectic twist map. From a criterion of total destruction of Lagrangian tori found by Herman and an approximation lemma, total destruction of Lagrangian tori under C^ω perturbations is completed.

2. DESTRUCTION OF A LAGRANGIAN TORUS

2.1. Case with 2 degrees of freedom

Based on the correspondence between exact area-preserving twist maps and Hamiltonian systems with 2 degrees of freedom, it is sufficient to consider the destruction of invariant circle for exact area-preserving twist map.

2.1.1. Preliminaries of exact area-preserving twist map

2.1.1.1. Minimal configuration

Let f: $\mathbb{T}\times\mathbb{R}\to\mathbb{T}\times\mathbb{R}$ ($\mathbb{T}=\mathbb{R}/\mathbb{Z}$) be an exact area-preserving monotone twist map and h: $\mathbb{R}^2\to\mathbb{R}^2$ be a generating function for the lift F of f to $\mathbb{R}^2$, namely F is generated by the following equations

$$\begin{cases} y=-\partial_1 h(x,x'),\\ y'=\partial_2 h(x,x'),\end{cases}$$

where $F(x,y)=(x',y')$. The lift F gives rise to a dynamical system whose orbits are given by the images of points of $\mathbb{R}^2$ under the successive iterates of F. The orbit of the point (x_0,y_0) is the bi-infinite sequence

$$\{...,(x_{-k},y_{-k}),...,(x_{-1},y_{-1}),(x_0,y_0),(x_1,y_1),...,(x_k,y_k),...\},$$

where $(x_k,y_k)=F(x_{k-1},y_{k-1})$. The sequence

$$(...,x_{-k},...,x_{-1},x_0,x_1,...,x_k,...)$$

denoted by $(x_i)_{i\in\mathbb{Z}}$ is called stationary configuration if it stratifies the identity

$$\partial_1 h(x_i,x_{i+1})+\partial_2 h(x_{i-1},x_i)=0,\ \text{for every } i\in\mathbb{Z}.$$

Given a sequence of points $(z_i,...,z_j)$, we can associate its action

$$h(z_i,...,z_j)=\sum_{i\leq s<j} h(z_s,z_{s+1}).$$

A configuration $(x_i)_{i\in\mathbb{Z}}$ is called minimal if for any $i<j\in\mathbb{Z}$, the segment of $(x_i,...,x_j)$ minimizes $h(z_i,...,z_j)$ among all segments $(z_i,...,z_j)$ of the configuration satisfying $z_i=x_i$ and $z_j=x_j$. It is easy to see that every minimal configuration is a stationary configuration. By [Ba], minimal configurations satisfy a group of remarkable properties as follows:

- Two distinct minimal configurations cross at most once, which is so called Aubry's crossing lemma.

- For every minimal configuration $\mathbf{x} = (x_i)_{i\in\mathbb{Z}}$, the limit
$$\rho(\mathbf{x}) = \lim_{n\to\infty} \frac{x_{i+n} - x_i}{n}$$
exists and doesn't depend on $i \in \mathbb{Z}$. $\rho(\mathbf{x})$ is called the rotation number of $\mathbf{x}$.
- For every $\omega \in \mathbb{R}$, there exists a minimal configuration with rotation number ω. Following the notations of [B], the set of all minimal configurations with rotation number ω is denoted by M^h_ω, which can be endowed with the topology induced from the product topology on $\mathbb{R}^\mathbb{Z}$. If $\mathbf{x} = (x_i)_{i\in\mathbb{Z}}$ is a minimal configuration, considering the projection $pr:\ M^h_\omega \to \mathbb{R}$ defined by $pr(\mathbf{x}) = x_0$, we set $\mathcal{A}^h_\omega = pr(M^h_\omega)$.
- If $\omega \in \mathbb{Q}$, say $\omega = p/q$ (in lowest terms), then it is convenient to define the rotation symbol to detect the structure of $M^h_{p/q}$. If $\mathbf{x}$ is a minimal configuration with rotation number p/q, then the rotation symbol $\sigma(\mathbf{x})$ of $\mathbf{x}$ is defined as follows
$$\sigma(\mathbf{x}) = \begin{cases} p/q+, & \text{if } x_{i+q} > x_i + p \text{ for all } i,\\ p/q, & \text{if } x_{i+q} = x_i + p \text{ for all } i,\\ p/q-, & \text{if } x_{i+q} < x_i + p \text{ for all } i. \end{cases}$$
Moreover, we set
$$M^h_{p/q^+} = \{\mathbf{x} \text{ a is minimal configuration with rotation symbol } p/q \text{ or } p/q+\},$$
$$M^h_{p/q^-} = \{\mathbf{x} \text{ a is minimal configuration with rotation symbol } p/q \text{ or } p/q-\},$$
then both M^h_{p/q^+} and M^h_{p/q^+} are totally ordered. Namely, every two configurations in each of them do not cross. We denote $pr(M^h_{p/q^+})$ and $pr(M^h_{p/q^-})$ by $\mathcal{A}^h_{p/q^+}$ and $\mathcal{A}^h_{p/q^-}$ respectively.
- If $\omega \in \mathbb{R}\backslash\mathbb{Q}$ and $\mathbf{x}$ is a minimal configuration with rotation number ω, then $\sigma(\mathbf{x}) = \omega$ and M^h_ω is totally ordered.
- $\mathcal{A}^h_\omega$ is a closed subset of $\mathbb{R}$ for every rotation symbol ω.

2.1.1.2. Peierls's barrier

In [Ma4], Mather introduced the notion of Peierls's barrier and gave a criterion of existence of invariant circle. Namely, the exact area-preserving monotone twist map generated by h admits an invariant circle with rotation number ω if and only if the Peierls's barrier $P^h_\omega(\xi)$ vanishes identically for all $\xi \in \mathbb{R}$. The Peierls's barrier is defined as follows:

- If $\xi \in \mathcal{A}^h_\omega$, we set $P^h_\omega(\xi)$=0.
- If $\xi \notin \mathcal{A}^h_\omega$, since $\mathcal{A}^h_\omega$ is a closed set in $\mathbb{R}$, then ξ is contained in some complementary interval (ξ^-, ξ^+) of $\mathcal{A}^h_\omega$ in $\mathbb{R}$. By the definition of $\mathcal{A}^h_\omega$, there exist minimal configurations with rotation symbol ω, $\mathbf{x}^- = (x^-_i)_{i\in\mathbb{Z}}$ and $\mathbf{x}^+ = (x^+_i)_{i\in\mathbb{Z}}$ satisfying $x^-_0 = \xi^-$ and $x^+_0 = \xi^+$. For every configuration $\mathbf{x} = (x_i)_{i\in\mathbb{Z}}$ satisfying $x^-_i \le x_i \le x^+_i$, we set
$$G_\omega(\mathbf{x}) = \sum_I (h(x_i, x_{i+1}) - h(x^-_i, x^-_{i+1})),$$

where $I = \mathbb{Z}$, if ω is not a rational number, and $I = \{0, ..., q-1\}$, if $\omega = p/q$. $P^h_\omega(\xi)$ is defined as the minimum of $G_\omega(\mathbf{x})$ over the configurations $\mathbf{x} \in \Pi = \prod_{i\in I}[x_i^-, x_i^+]$ satisfying $x_0 = \xi$. Namely

$$P^h_\omega(\xi) = \min_{\mathbf{x}}\{G_\omega(\mathbf{x})|\mathbf{x} \in \Pi \text{ and } \ x_0 = \xi\}.$$

By [Ma4], $P^h_\omega(\xi)$ is a non-negative periodic function of the variable $\xi \in \mathbb{R}$ with the modulus of continuity with respect to ω.

2.1.2. C^∞ case

2.1.2.1. Construction of the generating functions

Consider a completely integrable system with the generating function

$$h_0(x, x') = \frac{1}{2}(x - x')^2 \quad x, x' \in \mathbb{R}.$$

We construct the perturbation consisting of two parts. The first one is

$$u_n(x) = \frac{1}{n^a}(1 - \cos(2\pi x)) \quad x \in \mathbb{R}, \tag{2.1.1}$$

where $n \in \mathbb{N}$ and a is a positive constant independent of n. The second one is a non negative function $v_n(x)$ satisfying

$$\begin{cases} v_n(x+1) = v_n(x), \\ \operatorname{supp} v_n \cap [0,1] \subset [\frac{1}{2} - \frac{1}{n^a}, \frac{1}{2} + \frac{1}{n^a}], \\ \max v_n = n^{-s}, \\ ||v_n||_{C^k} \sim n^{-s'}, \end{cases} \tag{2.1.2}$$

where $f \sim g$ means that $\frac{1}{C}g < f < Cg$ holds for a constant $C > 1$. For further deduction, we need $s' > a$. It is enough to take $s = (k+2)a$ for achieving that. The generating function of the nearly integrable system is constructed as follow:

$$h_n(x, x') = h_0(x, x') + u_n(x') + v_n(x'), \tag{2.1.3}$$

where $n \in \mathbb{N}$. Moreover, we have the following theorem.

Theorem 2.1 *For $\omega \in \mathbb{R}\backslash\mathbb{Q}$ and n large enough, the exact area-preserving monotone twist map generated by h_n does not admit any invariant circles with the rotation number satisfying*

$$|\omega| < n^{-\frac{a}{2}-\delta},$$

where δ is a small positive constant independent of n.

We will prove Theorem 2.1 in the following sections. First of all, based on the theorem, we verify that our example has the property aforementioned in Section 1.

If $\omega \in \mathbb{Q}$, then the invariant circles with rotation number ω could be easily destructed by an analytic perturbation arbitrarily close to 0. Therefore it suffices to consider the irrational ω. The case with a given irrational rotation number can be easily reduced to the one with a small enough rotation number. More precisely,

Lemma 2.2 *Let* h_P *be a generating function as follow*

$$h_P(x,x')=h_0(x,x')+P(x'),$$

where P *is a periodic function of periodic* 1. *Let* $Q(x)=q^{-2}P(qx), q\in\mathbb{N}$*, then the exact area-preserving monotone twist map generated by* $h_Q(x,x')=h_0(x,x')+Q(x')$ *admits an invariant circle with rotation number* $\omega\in\mathbb{R}\backslash\mathbb{Q}$ *if and only if the exact area-preserving monotone twist map generated by* h_P *admits an invariant circle with rotation number* $q\omega-p, p\in\mathbb{Z}$.

We omit the proof and for more details, see [H2]. For the sake of simplicity of notations, we denote Q_{q_n} by Q_n and the same to u_{q_n}, v_{q_n} and h_{q_n}. Let

$$Q_n(x)={q_n}^{-2}(u_n(q_nx)+v_n(q_nx)),$$

where $(q_n)_{n\in\mathbb{N}}$ is a sequence satisfying Dirichlet approximation

$$|q_n\omega-p_n|<\frac{1}{q_n},\tag{2.1.4}$$

where $p_n\in\mathbb{Z}$ and $q_n\in\mathbb{N}$. Since $\omega\in\mathbb{R}\backslash\mathbb{Q}$, we say $q_n\to\infty$ as $n\to\infty$. Let $\tilde{h}_n(x,x')=h_0(x,x')+Q_n(x')$, we have

Corollary 2.3 *For a given rotation number* $\omega\in\mathbb{R}\backslash\mathbb{Q}$ *and every* ε*, there exists* N *such that for* $n>N$*, the exact area-preserving monotone map generated by* $\tilde{h}_n$ *admits no invariant circle with rotation number* ω *and*

$$||\tilde{h}_n-h_0||_{C^{4-\delta'}}<\varepsilon,$$

where δ' *is a small positive constant independent of* n.

Proof Based on Theorem 2.1 and Dirichlet approximation (2.1.54), it suffices to take

$$\frac{1}{q_n}\leq\frac{1}{{q_n}^{\frac{a}{2}+\delta}},$$

which implies

$$a\leq 2-2\delta.\tag{2.1.5}$$

From (2.1.30), (2.1.2) and (2.2.38), it follows that

$$\begin{aligned}||\tilde{h}_n(x,x')-h_0(x,x')||_{C^r}&\\ &=||Q_n(x')||_{C^r},\\ &\leq {q_n}^{-2}(||u_n(q_nx')||_{C^r}+||v_n(q_nx')||_{C^r}),\\ &\leq {q_n}^{-2}({q_n}^{-a}(2\pi)^r{q_n}^r+C_1{q_n}^{-s'}{q_n}^r),\\ &\leq C_2{q_n}^{r-a-2},\end{aligned}$$

where C_1, C_2 are positive constants only depending on r.

To complete the proof, it is enough to make $r-a-2<0$, which together with (2.1.55) implies

$$r<a+2\leq 4-2\delta.$$

We set $\delta'=2\delta$, then the proof of Corollary 2.3 is completed. □

The following sections are devoted to prove Theorem 2.1. For simplicity, we don't distinguish the constant C in following different estimate formulas.

2.1.2.2. Estimate of lower bound of $P_{0^+}^{h_n}$

In this section, we will estimate the lower bound of $P_{0^+}^{h_n}$ at a given point. To achieve that, we need to estimate the distances of pairwise adjacent elements of the minimal configuration.

2.1.2.2.1. A spacing lemma

Lemma 2.4 *Let $(x_i)_{i\in\mathbb{Z}}$ be a minimal configuration of $\bar{h}_n$ with rotation symbol 0^+, then*

$$x_{i+1}-x_i \geq C(n^{-\frac{a}{2}}), \quad \textit{for} \quad x_i \in \left[\frac{1}{4},\frac{3}{4}\right],$$

where $\bar{h}_n(x_i,x_{i+1}) = h_0(x_i,x_{i+1}) + u_n(x_{i+1})$.

Proof Without loss of generality, we assume $x_i \in [0,1]$ for all $i \in \mathbb{Z}$. By Aubry's crossing lemma, we have

$$0 < \ldots < x_{i-1} < x_i < x_{i+1} < \ldots < 1.$$

We consider the configuration $(\xi_i)_{i\in\mathbb{Z}}$ defined by

$$\xi_j = \begin{cases} x_j, & j < i, \\ x_{j+1}, & j \geq i. \end{cases}$$

Since $(x_i)_{i\in\mathbb{Z}}$ is minimal, we have

$$\sum_{i\in\mathbb{Z}} \bar{h}_n(\xi_i,\xi_{i+1}) - \sum_{i\in\mathbb{Z}} \bar{h}_n(x_i,x_{i+1}) \geq 0.$$

By the definitions of $\bar{h}_n$ and $(\xi_i)_{i\in\mathbb{Z}}$, we have

$$\begin{aligned} 0 &\leq \sum_{i\in\mathbb{Z}} \bar{h}_n(\xi_i,\xi_{i+1}) - \sum_{i\in\mathbb{Z}} \bar{h}_n(x_i,x_{i+1}) \\ &= \bar{h}_n(x_{i-1},x_{i+1}) - \bar{h}_n(x_{i-1},x_i) - \bar{h}_n(x_i,x_{i+1}) \\ &= (x_{i+1}-x_i)(x_i-x_{i-1}) - u_n(x_i). \end{aligned}$$

Moreover,

$$u_n(x_i) \leq (x_{i+1}-x_i)(x_i-x_{i-1}) \leq \frac{1}{4}(x_{i+1}-x_{i-1})^2.$$

Therefore,

$$x_{i+1}-x_{i-1} \geq 2\sqrt{u_n(x_i)}.$$

For $x_i \in [\frac{1}{4},\frac{3}{4}]$, $u_n(x_i) \geq n^{-a}$, hence,

$$x_{i+1}-x_{i-1} \geq 2n^{-\frac{a}{2}}. \tag{2.1.6}$$

Since $(x_i)_{i\in\mathbb{Z}}$ is a stationary configuration, we have

$$\begin{aligned} x_{i+1}-x_i &= -\partial_1\bar{h}_n(x_i,x_{i+1}), \\ &= \partial_2\bar{h}_n(x_{i-1},x_i), \\ &= x_i - x_{i-1} + u_n'(x_i). \end{aligned}$$

Since $u_n'(x) = \frac{2\pi}{n^a}\sin(2\pi x)$, it follows from (2.1.6) that

$$x_{i+1}-x_i \geq C(n^{-\frac{a}{2}}), \quad x_i \in \left[\frac{1}{4},\frac{3}{4}\right].$$

The proof of Lemma 2.4 is completed. $\square$

2.1.2.2.2. The lower bound of $P_{0^+}^{h_n}$

By the definition of v_n, $\operatorname{supp} v_n \cap [0,1] \subset [\frac{1}{2} - \frac{1}{n^a}, \frac{1}{2} + \frac{1}{n^a}]$ and $v_n(x+1) = v_n(x)$. Let $(x_i)_{i\in\mathbb{Z}}$ be the minimal configuration of $\bar{h}_n(x_i, x_{i+1}) = h_0(x_i, x_{i+1}) + u_n(x_{i+1})$ with rotation symbol 0^+ satisfying $x_0 = \frac{1}{2} - \frac{1}{n^a}$, then

$$(x_i)_{i\in\mathbb{Z}} \cap \operatorname{supp} v_n = \emptyset.$$

Moreover, for all $i \in \mathbb{Z}$,

$$v_n(x_i) = 0.$$

Let $(\xi_i)_{i\in\mathbb{Z}}$ be a minimal configuration of h_n defined by (2.2.38) with rotation symbol 0^+ satisfying $\xi_0 = \eta$, where η satisfies $v_n(\eta) = \max v_n(x) = n^{-s}$, then

$$\begin{aligned}
&\sum_{i\in\mathbb{Z}} (h_n(\xi_i, \xi_{i+1}) - h_n(\xi_i^-, \xi_{i+1}^-)) \\
&\qquad \geq v_n(\eta) + \sum_{i\in\mathbb{Z}} \bar{h}_n(\xi_i, \xi_{i+1}) - \sum_{i\in\mathbb{Z}} h_n(\xi_i^-, \xi_{i+1}^-), \\
&\qquad \geq v_n(\eta) + \sum_{i\in\mathbb{Z}} \bar{h}_n(x_i, x_{i+1}) - \sum_{i\in\mathbb{Z}} h_n(x_i, x_{i+1}), \\
&\qquad = v_n(\eta) - \sum_{i\in\mathbb{Z}} v_n(x_{i+1}), \\
&\qquad = v_n(\eta).
\end{aligned}$$

Therefore,

$$P_{0^+}^{h_n}(\eta) = \min_{x_0=\eta} \sum_{i\in\mathbb{Z}} (h_n(x_i, x_{i+1}) - h_n(x_i^-, x_{i+1}^-)) \geq v_n(\eta) = n^{-s}.$$

We conclude that there exists a point $\xi \in [\frac{1}{2} - \frac{1}{n^a}, \frac{1}{2} + \frac{1}{n^a}]$ such that

$$P_{0^+}^{h_n}(\xi) \geq n^{-s}. \tag{2.1.7}$$

2.1.2.3. The approximation from $P_{0^+}^{h_n}$ to $P_{\omega}^{h_n}$

In this section, we will prove the improvement of modulus of continuity of Peierls's barrier based on the hyperbolicity of h_n. Namely

Lemma 2.5 *For every irrational rotation symbol ω satisfying $0 < \omega < n^{-\frac{a}{2}-\delta}$, we have*

$$|P_{\omega}^{h_n}(\xi) - P_{0^+}^{h_n}(\xi)| \leq C \exp\left(-2n^{\frac{\delta}{2}}\right).$$

where $\xi \in \left[\frac{1}{2} - \frac{1}{n^a}, \frac{1}{2} + \frac{1}{n^a}\right]$ and δ is a small positive constant independent of n.

2.1.2.3.1. Some counting lemmas

To prove the lemma, we need to do some preliminary work. First of all, we count the number of the elements of a minimal configuration $(x_i)_{i\in\mathbb{Z}}$ with arbitrary rotation symbol ω in a given interval. With the method of [F], we can conclude the following lemma.

Lemma 2.6 *Let* $(x_i)_{i\in\mathbb{Z}}$ *be a minimal configuration of* h_n *with rotation symbol* $\omega > 0$*,* $J_n = \left[\exp\left(-n^{\frac{\delta}{2}}\right), \frac{1}{2}\right]$ *and* $\Lambda_n = \{i \in \mathbb{Z} |\, x_i \in J_n\}$*, then*

$$\sharp\Lambda_n \le Cn^{\frac{a}{2}+\frac{\delta}{2}},$$

where $\sharp\Lambda_n$ *denotes the number of elements in* Λ_n *and* δ *is a small positive constant independent of* n*.*

Proof Let $x^- = \exp\left(-n^{\frac{\delta}{2}}\right), x^+ = \frac{1}{2}$ and $\sigma = \left(\frac{x^+}{x^-}\right)^{\frac{1}{N}}$, hence,

$$\ln\sigma = \frac{\ln(x^+) - \ln(x^-)}{N}.$$

We choose $N \in \mathbb{N}$ such that $1 \le \ln\sigma \le 2$, then $N = \Omega\left(n^{\frac{\delta}{2}}\right)$.

We consider the partition of the interval $J_n = [x^-, x^+]$ into the subintervals $J_n^k = [\sigma^k x^-, \sigma^{k+1}x^-]$ where $0 \le k < N$. Hence, $J_n = \cup_{k=0}^{N-1} J_n^k$. We set $S_k = \{i \in \Lambda_n | (x_{i-1}, x_{i+1}) \subset J_n^k\}$ and $m_k = \sharp S_k$.

By the similar deduction as the one in Lemma 2.4, we have

$$x_{i+1} - x_{i-1} \ge 2\sqrt{u_n(x_i) + v_n(x_i)} \ge Cn^{-\frac{a}{2}}x_i, \quad \text{for} \quad x_i \in \left[0, \frac{1}{2}\right].$$

For simplicity of notation, we denote $Cn^{-\frac{a}{2}}$ by α_n.

If there exists k such that $i \in S_k$ for $(x_i)_{i\in\mathbb{Z}}$, then $x_{i+1} - x_{i-1} \ge \alpha_n\sigma^k x^-$, moreover,

$$m_k\alpha_n\sigma^k x^- \le 2\mathcal{L}(J_n^k) = 2(\sigma - 1)\sigma^k x^-,$$

where $\mathcal{L}(J_n^k)$ denotes the length of the interval of J_n^k. Hence $m_k \le 2(\sigma - 1)\alpha_n^{-1}$.

On the other hand, if $i \in \Lambda_n \backslash \cup_{k=0}^{N-1} S_k$, then there exists l satisfying $0 \le l < N$ such that

$$x_{i-1} < \sigma^l x^- < x_{i+1}.$$

Hence,

$$\sharp\{i \in \Lambda_n | i \notin S_k \text{ for any } k\} \le 2N.$$

Therefore,

$$\sharp(\Lambda_n) \le 2N(\sigma - 1)\alpha_n^{-1} + 2N.$$

Since $1 \le \ln\sigma \le 2$ and $N = \Omega\left(n^{\frac{\delta}{2}}\right)$, then we have

$$\sharp\Lambda_n \le Cn^{\frac{a}{2}+\frac{\delta}{2}}.$$

The proof of Lemma 2.6 is completed. □

Remark 2.7 *Let* $(x_i)_{i\in\mathbb{Z}}$ *be a minimal configuration of* h_n *defined by* (2.2.38) *with rotation symbol* $\omega > 0$*, An argument as similar as the one in Lemma 2.6 implies that*

$$\sharp\left\{i \in \mathbb{Z} \middle| x_i \in \left[\exp\left(-n^{\frac{\delta}{2}}\right), 1 - \exp\left(-n^{\frac{\delta}{2}}\right)\right]\right\} \le Cn^{\frac{a}{2}+\frac{\delta}{2}}.$$

It is easy to count the number of the elements of a minimal configuration with irrational rotation symbol. More precisely, we have the following lemma.

Lemma 2.8 *Let* $(x_i)_{i\in\mathbb{Z}}$ *be a minimal configuration with rotation number* $\omega \in \mathbb{R}\backslash\mathbb{Q}$*. Then for every interval* I_k *of length* k*,* $k \in \mathbb{N}$*,*

$$\frac{k}{\omega} - 1 \le \sharp\{i \in \mathbb{Z} | x_i \in I_k\} \le \frac{k}{\omega} + 1.$$

Proof For every minimal configuration $(x_i)_{i\in\mathbb{Z}}$ with rotation number ω, there exists an orientation-preserving circle homeomorphism ϕ such that $\rho(\Phi) = \omega$, where $\Phi : \mathbb{R} \to \mathbb{R}$ denotes a lift of ϕ. Since $\omega \in \mathbb{R}\backslash\mathbb{Q}$, thanks to [H1], ϕ has a unique invariant probability measure $\bar{\mu}$ on $\mathbb{T}$ such that $\int_x^{\Phi(x)} d\bar{\mu} = \omega$ for every $x \in \mathbb{R}$. We denote $\int_x^{\Phi(x)} d\bar{\mu}$ by $\mu(x, \Phi(x))$. In particular,

$$\mu(x_i, x_{i+1}) = \omega, \quad \text{for every } i \in \mathbb{Z}.$$

From $\mu(I_k) = k$, it follow that

$$\begin{aligned} &\omega(\sharp\{i \in \mathbb{Z} | x_i \in I_k\} - 1) \le k, \\ &\omega(\sharp\{i \in \mathbb{Z} | x_i \in I_k\} + 1) \ge k, \end{aligned}$$

which completes the proof of Lemma 2.8. □

Based on Lemma 2.6 and Lemma 2.8, if $0 < \omega < n^{-\frac{a}{2}-\delta}$ and ω is irrational, then

$$\sharp\{i \in \mathbb{Z} | x_i \in I_1\} \ge \frac{1}{\omega} - 1 \ge C_1 n^{\frac{a}{2}+\delta} > C_2 n^{\frac{a}{2}+\frac{\delta}{2}}, \tag{2.1.8}$$

where I_1 denotes the closed interval of length 1. Moreover, we have the following conclusion.

Lemma 2.9 *Let* $(x_i)_{i\in\mathbb{Z}}$ *be a minimal configuration of* h_n *defined by* (2.2.38) *with rotation symbol* $0 < \omega < n^{-\frac{a}{2}-\delta}$*, then there exist* $j^-, j^+ \in \mathbb{Z}$ *such that*

$$\begin{aligned} &0 < x_{j^- -1} < x_{j^-} < x_{j^- +1} \le \exp(-n^{\frac{\delta}{2}}), \\ &1 - \exp(-n^{\frac{\delta}{2}}) \le x_{j^+ -1} < x_{j^+} < x_{j^+ +1} < 1. \end{aligned}$$

Proof By contradiction, we assume that there exist at most two points of $(x_i)_{i\in\mathbb{Z}}$ in $[0, \exp(-n^{\frac{\delta}{2}})]$, say x_m and x_{m+1}. It follows that $x_{m-1} < 0$ and $x_{m+2} > \exp(-n^{\frac{\delta}{2}})$. Hence, among the intervals $[x_{m-1}, x_m]$, $[x_m, x_{m+1}]$ and $[x_{m+1}, x_{m+2}]$, there exists at least one such that its length is not less than $\frac{1}{3}\exp(-n^{\frac{\delta}{2}})$. Without loss of generality, say $[x_{m+1}, x_{m+2}]$.

Since $(x_i)_{i\in\mathbb{Z}}$ is a stationary configuration, we have

$$x_{m+2} - x_{m+1} = x_{m+1} - x_m + u_n'(x_{m+1}),$$

where $u_n'(x_{m+1}) = \frac{2\pi}{n^a}\sin(2\pi x_{m+1})$. From $x_{m+1} \in [-\exp(-n^{\frac{\delta}{2}}), \exp(-n^{\frac{\delta}{2}})]$, it follows that

$$|u_n'(x_{m+1})| \le Cn^{-a}\exp(-n^{\frac{\delta}{2}}),$$

which implies there exists N independent of n such that $[-\exp(-n^{\frac{\delta}{2}}), \exp(-n^{\frac{\delta}{2}})]$ contains at most N points of $(x_i)_{i\in\mathbb{Z}}$.

On the other hand, by (2.1.8), we have that for n large enough, the number of points of $(x_i)_{i\in\mathbb{Z}}$ in $[-\exp(-n^{\frac{\delta}{2}}), \exp(-n^{\frac{\delta}{2}})]$ is also large enough, which is a contradiction. Therefore, there exists $j^- \in \mathbb{Z}$ such that

$$0 \leq x_{j^- -1} < x_{j^-} < x_{j^- +1} < \exp(-n^{\frac{\delta}{2}}).$$

Similarly, there exists $j^+ \in \mathbb{Z}$ such that

$$1 - \exp(-n^{\frac{\delta}{2}}) \leq x_{j^+ -1} < x_{j^+} < x_{j^+ +1} < 1.$$

The proof of Lemma 2.9 is completed. □

Remark 2.10 *From the proof of Lemma 2.9, it is easy to see that each of* $[0, \exp(-n^{\frac{\delta}{2}})]$ *and* $[1-\exp(-n^{\frac{\delta}{2}}), 1]$ *contains a large number of points of the minimal configuration* $(x_i)_{i\in\mathbb{Z}}$ *for* n *large enough.*

By Lemma 2.8 and Lemma 2.9, without loss of generality, one can assume that

$$j^+ - j^- \geq C\left(n^{\frac{a}{2}+\frac{2\delta}{3}}\right). \tag{2.1.9}$$

If $\xi \in \mathcal{A}_\omega^{h_n}$, then $P_\omega^{h_n}(\xi) = 0$. Hence, it suffices to consider the case with $\xi \notin \mathcal{A}_\omega^{h_n}$ for destruction of invariant circles. Let (ξ^-, ξ^+) be the complementary interval of $\mathcal{A}_\omega^{h_n}$ in $\mathbb{R}$ and contains ξ. Let $\boldsymbol{\xi}^{\pm} = (\xi_i^{\pm})_{i\in\mathbb{Z}}$ be the minimal configurations with rotation symbol ω satisfying $\xi_0^{\pm} = \xi^{\pm}$ and let $(\xi_i)_{i\in\mathbb{Z}}$ be a minimal configuration of h_n with rotation symbol ω satisfying $\xi_0 = \xi$ and $\xi_i^- \leq \xi_i \leq \xi_i^+$. By the definition of Peierls barrier, we have

$$P_\omega^{h_n}(\xi) = \sum_{i\in\mathbb{Z}} (h_n(\xi_i, \xi_{i+1}) - h_n(\xi_i^-, \xi_{i+1}^-)).$$

Since $P_\omega^{h_n}(\xi)$ is 1-periodic with respect to ξ, without loss of generality, we assume that $\xi \in [0,1]$. We set $d(x) = \min\{|x|, |x-1|\}$ and denote $\exp(-n^{\frac{\delta}{2}})$ by $\epsilon(n)$. By Lemma 2.9, there exist i^-, i^+ such that

$$d(\xi_i^-) < \epsilon(n) \quad \text{and} \quad \xi_{i+1}^- - \xi_{i-1}^- \leq \epsilon(n) \quad \text{for} \quad i = i^-,\ i^+. \tag{2.1.10}$$

Thanks to Aubry's crossing lemma, we have $\xi_i^- \leq \xi_i \leq \xi_i^+ \leq \xi_{i+1}^-$. Hence,

$$\xi_i - \xi_i^- \leq \epsilon(n) \quad \text{for} \quad i = i^-,\ i^+.$$

2.1.2.3.2. Proof of lemma 2.5

In the following, we will prove Lemma 2.5 with the method similar to the one developed by Mather in [M3]. The proof can be proceeded in the following two steps.

Step 1 We consider the number of the elements in a segment of the configuration as the length of the segment. In the first step, we approximate $P_\omega^{h_n}(\xi)$ for $\xi \in$

$\left[\frac{1}{2}-\frac{1}{n^a},\frac{1}{2}+\frac{1}{n^a}\right]$ by the difference of the actions of the segments of length i^+-i^-+1. To achieve that, we define the following configurations

$$x_i=\begin{cases}\xi_i, & i\neq i^-,\ i^+,\\ \xi_i^-, & i=i^-,\ i^+,\end{cases}\qquad \text{and}\qquad y_i=\begin{cases}\xi_i, & i^-<i<i^+,\\ \xi_i^-, & i\leq i^-,\ i\geq i^+,\end{cases}$$

where $(\xi_i)_{i\in\mathbb{Z}}$ is a minimal configuration. It is easy to see that $\xi_0=\xi$ is contained both of $(x_i)_{i\in\mathbb{Z}}$ and $(y_i)_{i\in\mathbb{Z}}$ up to the rearrangement of the index i since $\xi\in\left[\frac{1}{2}-\frac{1}{n^a},\frac{1}{2}+\frac{1}{n^a}\right]$. Hence, by the minimality of $(\xi_i)_{i\in\mathbb{Z}}$ satisfying $\xi_0=\xi$, we have

$$P_\omega^{h_n}(\xi)\leq\sum_{i\in\mathbb{Z}}(h_n(y_i,y_{i+1})-h_n(\xi_i^-,\xi_{i+1}^-)). \tag{2.1.11}$$

Since ω is irrational, then $(x_i)_{i\in\mathbb{Z}}$ is asymptotic to $(\xi_i^-)_{i\in\mathbb{Z}}$, which together with the minimality of $(\xi_i^-)_{i\in\mathbb{Z}}$ yields

$$\sum_{i\in\mathbb{Z}}(h_n(y_i,y_{i+1})-h_n(\xi_i^-,\xi_{i+1}^-)\leq\sum_{i\in\mathbb{Z}}(h_n(x_i,x_{i+1})-h_n(\xi_i^-,\xi_{i+1}^-)). \tag{2.1.12}$$

We set

$$h(x_i,...,x_j)=\sum_{i\leq s<j}h(x_s,x_{s+1}),$$

then

$$\sum_{i\in\mathbb{Z}}(h_n(x_i,x_{i+1})-h_n(\xi_i,\xi_{i+1}))=\sum_{i=i^-,i^+}(h_n(\xi_{i-1},\xi_i^-,\xi_{i+1})-h_n(\xi_{i-1},\xi_i,\xi_{i+1})).$$

By the construction of v_n and Lemma 2.9, we have $v_n(\xi_{i^-}),\ v_n(\xi_{i^-}^-)=0$. It follows that

$$\begin{aligned}
&h_n(\xi_{i^--1},\xi_{i^-}^-,\xi_{i^-+1})-h_n(\xi_{i^--1},\xi_{i^-},\xi_{i^-+1})\\
&\quad=h_n(\xi_{i^--1},\xi_{i^-}^-)+h_n(\xi_{i^-}^-,\xi_{i^-+1})-h_n(\xi_{i^--1},\xi_{i^-})-h_n(\xi_{i^-},\xi_{i^-+1}),\\
&\quad=(\xi_{i^-}-\xi_{i^-}^-)(\xi_{i^--1}+\xi_{i^-}^-+\xi_{i^-}+\xi_{i^-+1})+u_n(\xi_{i^-})-u_n(\xi_{i^-}^-),\\
&\quad\leq 4(\xi_{i^-}-\xi_{i^-}^-)\epsilon(n)+u_n'(\eta)(\xi_{i^-}-\xi_{i^-}^-),\\
&\quad\leq 4\epsilon(n)^2+\frac{2\pi}{n^a}\sin(2\pi\eta)\epsilon(n),\\
&\quad\leq C\epsilon(n)^2,
\end{aligned}$$

where $\eta\in(\xi_{i^-},\xi_{i^-}^-)$. It is similar to obtain

$$h_n(\xi_{i^+-1},\xi_{i^+}^-,\xi_{i^++1})-h_n(\xi_{i^+-1},\xi_{i^+},\xi_{i^++1})\leq C\epsilon(n)^2.$$

Hence,

$$\sum_{i\in\mathbb{Z}}(h_n(x_i,x_{i+1})-h_n(\xi_i,\xi_{i+1}))\leq C\epsilon(n)^2. \tag{2.1.13}$$

Moreover,

$$\begin{aligned}
&\sum_{i\in\mathbb{Z}}(h_n(x_i,x_{i+1})-h_n(\xi_i^-,\xi_{i+1}^-))\\
&\qquad=\sum_{i\in\mathbb{Z}}(h_n(x_i,x_{i+1})-h_n(\xi_i,\xi_{i+1})+h_n(\xi_i,\xi_{i+1})-h_n(\xi_i^-,\xi_{i+1}^-)),\\
&\qquad=\sum_{i\in\mathbb{Z}}(h_n(x_i,x_{i+1})-h_n(\xi_i,\xi_{i+1}))+P_\omega^{h_n}(\xi),\\
&\qquad\leq P_\omega^{h_n}(\xi)+C\epsilon(n)^2.
\end{aligned}$$

Therefore, it follows from (2.1.11) and (2.1.12) that

$$P_\omega^{h_n}(\xi) \le \sum_{i\in\mathbb{Z}} (h_n(y_i, y_{i+1}) - h_n(\xi_i^-, \xi_{i+1}^-)) \le P_\omega^{h_n}(\xi) + C\epsilon(n)^2, \tag{2.1.14}$$

where

$$\sum_{i\in\mathbb{Z}} (h_n(y_i, y_{i+1}) - h_n(\xi_i^-, \xi_{i+1}^-)) = h_n(y_{i^-}, ..., y_{i^+}) - h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-). \tag{2.1.15}$$

Step 2 It follows from [M4] that the Peierls's barrier $P_{0^+}^{h_n}(\xi)$ could be defined as follows

$$P_{0^+}^{h_n}(\xi) = \min_{\eta_0=\xi} \sum_{i\in\mathbb{Z}} h_n(\eta_i, \eta_{i+1}) - \min \sum_{i\in\mathbb{Z}} h_n(z_i, z_{i+1}), \tag{2.1.16}$$

where $(\eta_i)_{i\in\mathbb{Z}}$ and $(z_i)_{i\in\mathbb{Z}}$ are monotone increasing configurations limiting on 0, 1. We set

$$\begin{cases} K(\xi) = \min_{\eta_0=\xi} \sum_{i\in\mathbb{Z}} h_n(\eta_i, \eta_{i+1}), \\ K = \min \sum_{i\in\mathbb{Z}} h_n(z_i, z_{i+1}). \end{cases}$$

First of all, it is easy to see that $K(\xi)$ and K are bounded. Second, $P_{0^+}^{h_n}(\xi) = 0$ for $\xi = 0$ or 1. Hence, we only need to consider the case with $\xi \in (0, 1)$. Following the ideas of [M6], let $\boldsymbol{\xi}^-$ and $\boldsymbol{\xi}^+$ be minimal configurations of rotation symbol 0^+ and let (ξ_0^-, ξ_0^+) be the complementary interval of $\mathcal{A}_{0^+}^{h_n}$ and contains ξ. Based on the definition

$$P_{0^+}^{h_n}(\xi) = \min_{x_0=\xi} \{G_{0^+}(\mathbf{x}) | \xi_i^- \le \zeta_i \le \xi_i^+\},$$

where

$$G_{0^+}(\boldsymbol{\zeta}) = \sum_{i\in\mathbb{Z}} (h_n(\zeta_i, \zeta_{i+1}) - h_n(\xi_i^-, \xi_{i+1}^-)) = -K + \sum_{i\in\mathbb{Z}} h_n(\zeta_i, \zeta_{i+1}),$$

the proof of (2.1.16) will be completed when we verify that the configuration $(\zeta_i)_{i\in\mathbb{Z}}$ achieving the minimum in the definition of $K(\xi)$ satisfies $\xi_i^- \le \zeta_i \le \xi_i^+$. It can be easily obtained by Aubry's crossing lemma. In fact, since $(\xi_i^-)_{i\in\mathbb{Z}}$ and $(\zeta_i)_{i\in\mathbb{Z}}$ are minimal and both are α-asymptotic to 0 as well as ω-asymptotic to 1, by Aubry's crossing lemma, $(\xi_i^-)_{i\in\mathbb{Z}}$ and $(\zeta_i)_{i\in\mathbb{Z}}$ do not cross. Similarly $(\xi_i^+)_{i\in\mathbb{Z}}$ and $(\zeta_i)_{i\in\mathbb{Z}}$ do not cross. It follows from $\zeta_0 \in (\xi_0^-, \xi_0^+)$ that $(\zeta_i)_{i\in\mathbb{Z}}$ achieving the minimum in the definition of $K(\xi)$ satisfies $\xi_i^- \le \zeta_i \le \xi_i^+$.

In the following, we will compare K, $K(\xi)$ with $h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-)$, $h_n(y_{i^-}, ..., y_{i^+})$ respectively, here $h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-)$ and $h_n(y_{i^-}, ..., y_{i^+})$ are as the same as the notations in (2.1.15).

First, we consider K and $h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-)$. Let $(z_i)_{i\in\mathbb{Z}}$ be a monotone increasing configuration limiting on 0, 1 such that $K = \sum_{i\in\mathbb{Z}} h_n(z_i, z_{i+1})$. By Lemma 2.6,

$$\sharp\{i \in \mathbb{Z} | z_i \in [\epsilon(n), 1 - \epsilon(n)]\} \le Cn^{\frac{a}{2}+\frac{\delta}{2}}.$$

On the other hand, since $(z_i)_{i\in\mathbb{Z}}$ has the rotation number 0^+, then from (2.1.9), it follows that up to the rearrangement of the index i, there exists a subset of length $i^+ - i^-$ of $(z_i)_{i\in\mathbb{Z}}$, denoted by $\{z_{i^-}, z_{i^-+1}, \ldots, z_{i^+-1}, z_{i^+}\}$ such that

$$z_{i^-+1} \le \epsilon(n), \quad z_{i^+-1} \ge 1 - \epsilon(n).$$

By Lemma 2.9, $\xi^-_{i^-+1} > 0$ and $\xi^-_{i^+-1} < 1$ for the minimal configuration $(\xi^-_i)_{i\in\mathbb{Z}}$. We consider the configuration $(\bar{x}_i)_{i\in\mathbb{Z}}$ defined by

$$\begin{cases} \bar{x}_i = \xi^-_i, & i^- < i < i^+, \\ \bar{x}_i = 0, & i \le i^-, \\ \bar{x}_i = 1, & i \ge i^+. \end{cases}$$

By the definition of h_n, $h_n(\bar{x}_i, \bar{x}_{i+1}) = 0$ for $i < i^-$ or $i \ge i^+$, then

$$\sum_{i\in\mathbb{Z}} h_n(\bar{x}_i, \bar{x}_{i+1}) = h_n(\bar{x}_{i^-}, ..., \bar{x}_{i^+}).$$

Moreover, by the minimality of $(z_i)_{i\in\mathbb{Z}}$, we have

$$K \le \sum_{i\in\mathbb{Z}} h_n(\bar{x}_i, \bar{x}_{i+1}) = h_n(\bar{x}_{i^-}, ..., \bar{x}_{i^+}). \tag{2.1.17}$$

By the construction of h_n, we have $v_n(\bar{x}_{i^-+1}) = v_n(\xi^-_{i^-+1}) = 0$. Hence,

$$\begin{aligned} &h_n(\bar{x}_{i^-}, \bar{x}_{i^-+1}) - h_n(\xi^-_{i^-}, \xi^-_{i^-+1}) \\ &\quad= \frac{1}{2}(\bar{x}_{i^-} - \bar{x}_{i^-+1})^2 + u_n(\bar{x}_{i^-+1}) - \frac{1}{2}(\xi^-_{i^-} - \xi^-_{i^-+1})^2 - u_n(\xi^-_{i^-+1}), \\ &\quad= \frac{1}{2}(\xi^-_{i^-+1})^2 - \frac{1}{2}(\xi^-_{i^-+1} - \xi^-_{i^-})^2, \\ &\quad= \frac{1}{2}\xi^-_{i^-}(2\xi^-_{i^-+1} - \xi^-_{i^-}), \\ &\quad\le C\epsilon(n)^2. \end{aligned} \tag{2.1.18}$$

It is similar to obtain

$$h_n(\bar{x}_{i^+-1}, \bar{x}_{i^+}) - h_n(\xi^-_{i^+-1}, \xi^-_{i^+}) \le C\epsilon(n)^2. \tag{2.1.19}$$

Since

$$\begin{aligned} &h_n(\bar{x}_{i^-}, ..., \bar{x}_{i^+}) - h_n(\xi^-_{i^-}, ..., \xi^-_{i^+}) \\ &\quad= h_n(\bar{x}_{i^-}, \bar{x}_{i^-+1}) - h_n(\xi^-_{i^-}, \xi^-_{i^-+1}) + h_n(\bar{x}_{i^+-1}, \bar{x}_{i^+}) - h_n(\xi^-_{i^+-1}, \xi^-_{i^+}), \end{aligned}$$

then

$$h_n(\bar{x}_{i^-}, ..., \bar{x}_{i^+}) - h_n(\xi^-_{i^-}, ..., \xi^-_{i^+}) \le C\epsilon(n)^2. \tag{2.1.20}$$

From (2.1.17) and (2.1.20) we have

$$K \le h_n(\xi^-_{i^-}, ..., \xi^-_{i^+}) + C\epsilon(n)^2. \tag{2.1.21}$$

To obtain the reverse inequality of (2.2.45), we consider the configuration as follows

$$\begin{cases} \tilde{x}_i = z_i, & i^- < i < i^+, \\ \tilde{x}_i = 0, & i \le i^-, \\ \tilde{x}_i = 1, & i \ge i^+. \end{cases}$$

From the definition of h_n, it follows that $v_n(z_{i^-+1}) = 0$ and $h_n(z_i, z_{i+1}) \geq 0$ for all $i \in \mathbb{Z}$. Moreover, we have

$$\begin{aligned}
&h_n(\tilde{x}_{i^-}, ..., \tilde{x}_{i^+}) - K\\
&\quad = h_n(\tilde{x}_{i^-}, \tilde{x}_{i^-+1}) + h_n(\tilde{x}_{i^+-1}, \tilde{x}_{i^+}) - \sum_{i<i^-, i\geq i^+} h_n(z_i, z_{i+1}),\\
&\quad \leq \frac{1}{2}(z_{i^-+1})^2 + u_n(z_{i^-+1}) + \frac{1}{2}(z_{i^+-1} - 1)^2,\\
&\quad \leq u_n'(\eta) z_{i^-+1} + C_1\epsilon(n)^2,\\
&\quad \leq 2\pi n^{-a}\sin(2\pi\eta) z_{i^-+1} + C_1\epsilon(n)^2,\\
&\quad \leq C_2 n^{-a}(z_{i^-+1})^2 + C_1\epsilon(n)^2,\\
&\quad \leq C\epsilon(n)^2.
\end{aligned}$$

where $\eta \in (0, z_{i^-+1})$. Namely

$$h_n(\tilde{x}_{i^-}, ..., \tilde{x}_{i^+}) \leq K + C\epsilon(n)^2. \tag{2.1.22}$$

Furthermore, we consider the finite segment of the configuration defined by

$$\begin{cases} \eta_i = \tilde{x}_i, & i^- < i < i^+,\\ \eta_i = \xi_i^-, & i = i^-,\\ \eta_i = \xi_i^-, & i = i^+. \end{cases}$$

Then, the minimality of $(\xi_i^-)_{i\in\mathbb{Z}}$ implies $h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-) \leq h_n(\eta_{i^-}, ..., \eta_{i^+})$. Hence, by (2.1.22), we have

$$h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-) \leq K + C\epsilon(n)^2 + h_n(\eta_{i^-}, ..., \eta_{i^+}) - h_n(\tilde{x}_{i^-}, ..., \tilde{x}_{i^+}), \tag{2.1.23}$$

where

$$\begin{aligned}
&h_n(\eta_{i^-}, ..., \eta_{i^+}) - h_n(\tilde{x}_{i^-}, ..., \tilde{x}_{i^+})\\
&\quad = h_n(\eta_{i^-}, \eta_{i^-+1}) - h_n(\tilde{x}_{i^-}, \tilde{x}_{i^-+1}) + h_n(\eta_{i^+-1}, \eta_{i^+}) - h_n(\tilde{x}_{i^+-1}, \tilde{x}_{i^+}).
\end{aligned}$$

By the deduction as similar as (2.1.18), we have

$$\begin{aligned}
h_n(\eta_{i^-}, \eta_{i^-+1}) - h_n(\tilde{x}_{i^-}, \tilde{x}_{i^-+1}) \leq C\epsilon(n)^2,\\
h_n(\eta_{i^+-1}, \eta_{i^+}) - h_n(\tilde{x}_{i^+-1}, \tilde{x}_{i^+}) \leq C\epsilon(n)^2.
\end{aligned}$$

Moreover,

$$h_n(\eta_{i^-}, ..., \eta_{i^+}) - h_n(\tilde{x}_{i^-}, ..., \tilde{x}_{i^+}) \leq C\epsilon(n)^2. \tag{2.1.24}$$

Hence, from (2.1.23) and (2.13), it follows that

$$h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-) \leq K + C\epsilon(n)^2, \tag{2.1.25}$$

which together with (2.2.45) implies

$$|h_n(\xi_{i^-}^-, ..., \xi_{i^+}^-) - K| \leq C\epsilon(n)^2. \tag{2.1.26}$$

Next, we compare $h_n(y_{i^-}, ..., y_{i^+})$ with $K(\xi)$.

Since $(\xi_i)_{i\in\mathbb{Z}}$ is minimal among all configurations with rotation symbol ω satisfying $\xi_0=\xi$. By (2.1.10) and Aubry's crossing lemma, we have

$$d(\xi_i)\leq\epsilon(n),\quad \text{for } i=i^-,i^+,$$

where $d(\xi_i)=\min\{|\xi_i|,|\xi_i-1|\}$. By an argument as similar as the one in the comparison between K and $h_n(\xi^-_{i^-},...,\xi^-_{i^+})$, we have

$$|h_n(\xi_{i^-},...,\xi_{i^+})-K(\xi)|\leq C\epsilon(n)^2. \tag{2.1.27}$$

By the construction of $(y_i)_{i\in\mathbb{Z}}$, namely

$$y_i=\begin{cases}\xi_i, & i^-<i<i^+,\\ \xi_i^-, & i\leq i^-,\ i\geq i^+,\end{cases}$$

we have

$$\begin{aligned}&h_n(y_{i^-},...,y_{i^+})-h_n(\xi_{i^-},...,\xi_{i^+})\\ &\quad=h_n(\xi^-_{i^-},\xi_{i^-+1})-h_n(\xi_{i^-},\xi_{i^-+1})+h_n(\xi_{i^+-1},\xi^-_{i^+})-h_n(\xi_{i^+-1},\xi_{i^+}).\end{aligned}$$

By the deduction as similar as (2.1.20), we have

$$|h_n(y_{i^-},...,y_{i^+})-h_n(\xi_{i^-},...,\xi_{i^+})|\leq C\epsilon(n)^2. \tag{2.1.28}$$

Finally, from (2.1.14), (2.1.26), (2.1.27) and (2.1.28) we obtain

$$\begin{aligned}|P^{h_n}_{\omega}(\xi)-P^{h_n}_{0^+}(\xi)|&\leq|h_n(y_{i^-},...,y_{i^+})-h_n(\xi^-_{i^-},...,\xi^-_{i^+})+K-K(\xi)|+C_1\epsilon(n)^2,\\ &\leq|h_n(\xi_{i^-},...,\xi_{i^+})-K(\xi)|+|h_n(\xi^-_{i^-},...,\xi^-_{i^+})-K|\\ &\quad+|h_n(y_{i^-},...,y_{i^+})-h_n(\xi_{i^-},...,\xi_{i^+})|+C_1\epsilon(n)^2,\\ &\leq C\epsilon(n)^2,\\ &=C\exp\left(-2n^{\frac{\delta}{2}}\right),\end{aligned}$$

which completes the proof of Lemma 2.5. □

2.1.2.4. Proof of Theorem 2.1

Based on the preparation above, it is easy to prove Theorem 2.1. We assume that there exists an invariant circle with rotation number $0<\omega<n^{-\frac{a}{2}-\delta}$ for h_n, then $P^{h_n}_{\omega}(\xi)\equiv 0$ for every $\xi\in\mathbb{R}$. By Lemma 2.5, we have

$$|P^{h_n}_{0^+}(\xi)|\leq C\exp\left(-2n^{\frac{\delta}{2}}\right),\quad \text{for}\quad \xi\in\left[\frac{1}{2}-\frac{1}{n^a},\frac{1}{2}+\frac{1}{n^a}\right]. \tag{2.1.29}$$

On the other hand, (2.1.7) implies that there exists a point $\tilde{\xi}\in\left[\frac{1}{2}-\frac{1}{n^a},\frac{1}{2}+\frac{1}{n^a}\right]$ such that

$$P^{h_n}_{0^+}(\tilde{\xi})\geq n^{-s}.$$

Hence, we have

$$n^{-s}\leq C\exp\left(-2n^{\frac{\delta}{2}}\right).$$

It is an obvious contradiction for n large enough. Therefore, there exists no invariant circle with rotation number $0<\omega<n^{-\frac{a}{2}-\delta}$.

For $-n^{-\frac{a}{2}-\delta}<\omega<0$, by comparing $P^{h_n}_{\omega}(\xi)$ with $P^{h_n}_{0^-}(\xi)$, the proof is similar. We omit the details. Therefore, the proof of Theorem 2.1 is completed. □

2.1.3. C^ω case

We will prove the following theorem:

Theorem 2.11 *Given an integrable generating function h_0, a rotation number ω and a small positive constant δ, there exists a sequence of real-analytic $(h_n)_{n\in\mathbb{N}}$ such that $h_n \to h_0$ in the $C^{3-\delta}$ topology and the exact monotone area-preserving twist maps generated by $(h_n)_{n\in\mathbb{N}}$ admit no invariant circles with the rotation number ω.*

2.1.3.1. Construction of the generating functions

Consider a completely integrable system with the generating function

$$h_0(x,x') = \frac{1}{2}(x-x')^2, \quad x,x' \in \mathbb{R}.$$

We construct the perturbation consisting of two parts. The first one is

$$u_n(x) = \frac{1}{n^a}(1-\cos(2\pi x)), \quad x \in \mathbb{R}, \tag{2.1.30}$$

where $n \in \mathbb{N}$ and a is a positive constant independent of n.

We construct the second part of the perturbation in the following. Let $p_N(x)$ be a trigonometric polynomial of degree N. It is easy to see that for any $r > 0$,

$$||p_N(x)||_r \leq e^{rN}||p_N(x)||, \tag{2.1.31}$$

where $||p_N(x)||_r$ denotes the maximum of $|p_N(z)|$ in the strip $S_r = \{z \in \mathbb{C}|\, |\mathrm{Im} z| \leq r\}$ of width $2r$ in the complex plane and $||p_N(x)||$ denotes the maximum of $|p_N(x)|$ on the real line. Without loss of generality, we take $r = 1$, a.e.

$$||p_N(x)||_1 \leq e^N \max|p_N(x)|. \tag{2.1.32}$$

Then, by Cauchy estimates, for any fixed $s > 0$, we have

$$||p_N(x)||_{C^s} \leq C_s e^N \max|p_N(x)|, \tag{2.1.33}$$

where C_s is a constant depending only on s.

Based on Lemma 2.4, we need to construct a real analytic function with a "bump" in correspondence with the interval Λ_n satisfying

$$\mathcal{L}(\Lambda_n) \sim n^{-\frac{a}{2}} \quad \text{and} \quad \Lambda_n \subset \left[\frac{1}{4}, \frac{3}{4}\right], \tag{2.1.34}$$

where $\mathcal{L}(\Lambda_n)$ denotes the Lebesgue measure of Λ_n and $f \sim g$ means that $\frac{1}{C}g < f < Cg$ holds for some constant $C > 1$.

The "bump" will be accomplished by using Jackson's approximation theorem (see [Z, p115]). It states that let $\phi(x)$ be an k-times differentiable periodic function on $\mathbb{R}$, then for every $N \in \mathbb{N}$, there exists a trigonometric polynomial $p_N(x)$ of degree N such that

$$\max|p_N(x) - \phi(x)| \leq A_k N^{-k}||\phi(x)||_{C^k},$$

where A_k is a constant depending only on $k \in \mathbb{N}$.

We take a C^∞ bump function ϕ supported on the interval Λ_n, whose maximum is equal to 2. By (2.1.34), the length of Λ_n is bounded by $Cn^{-\frac{a}{2}}$, one can choose $\phi(x)$ such that

$$||\phi(x)||_{C^k} \sim \left(n^{\frac{a}{2}}\right)^k = n^{\frac{ak}{2}}. \tag{2.1.35}$$

Then, choosing N large enough to achieve

$$A_k N^{-k} ||\phi(x)||_{C^k} < \sigma \ll 1, \tag{2.1.36}$$

where σ will be determined in the following, by Jackson's approximation theorem, we can construct a trigonometric polynomial $p_N(x)$ of degree N such that:

$$\begin{cases} \max p_N(x) \geq 1, & \text{attained on } \Lambda_n, \\ |p_N(x)| \leq \sigma, & \text{on } [0,1]\backslash\Lambda_n. \end{cases} \tag{2.1.37}$$

By (3.4.2), we have

$$N \geq C\sigma^{-\frac{1}{k}} n^{\frac{a}{2}}. \tag{2.1.38}$$

Finally, we consider the normalized trigonometric polynomial

$$\tilde{p}_N(x) = e^{-2N}\left(\frac{p_N(x)}{\max p_N(x)}\right)^2. \tag{2.1.39}$$

From (2.1.33), $\tilde{p}_N(x)$ satisfies:

$$\begin{cases} \tilde{p}_N(x) \geq 0, \\ ||\tilde{p}_N(x)||_{C^s} \leq C, \\ \max \tilde{p}_N(x) = e^{-2N}, & \text{attained on } \Lambda_n, \\ |\tilde{p}_N(x)| \leq \sigma^2 e^{-2N}, & \text{on } [0,1]\backslash\Lambda_n. \end{cases} \tag{2.1.40}$$

Based on preparations above, we can construct the second part of the perturbation as follow

$$v_n(x) = u_n(x)\tilde{p}_N(x) = \frac{1}{n^a}(1-\cos 2\pi x)\tilde{p}_N(x). \tag{2.1.41}$$

It is easy to see v_n satisfies the following properties:

$$\begin{cases} v_n(x) \geq 0, \\ ||v_n(x)||_{C^s} \leq Cn^{-a}, \\ \max v_n(x) \geq e^{-2N} n^{-a}, & \text{attained on } \Lambda_n, \\ |v_n(x)| \leq C\sigma^2 e^{-2N} n^{-a}, & \text{on } [0,1]\backslash\Lambda_n. \end{cases} \tag{2.1.42}$$

So far, we complete the construction of the generating function of the nearly integrable system, a.e.

$$h_n(x,x') = h_0(x,x') + u_n(x') + v_n(x'), \tag{2.1.43}$$

where $n \in \mathbb{N}$.

2.1.3.2. Proof of Theorem 2.11

If $\omega \in \mathbb{Q}$, then the invariant circles with rotation number ω could be easily destructed by an analytic perturbation arbitrarily close to 0. Therefore it suffices to consider the irrational ω. Firstly, we prove the non-existence of invariant circles with a small enough rotation number. More precisely, we have the following Lemma:

Lemma 2.12 *For $\omega \in \mathbb{R}\backslash\mathbb{Q}$ and n large enough, the exact area-preserving monotone twist map generated by h_n admits no invariant circle with the rotation number satisfying*

$$|\omega| < n^{-a-\delta},$$

where δ is a small positive constant independent of n.

Proof First of all, we estimate the lower bound of $P^{h_n}_{0^+}$. Let $(\xi_i)_{i\in\mathbb{Z}}$ be a minimal configuration of h_n defined by (2.2.38) with rotation symbol 0^+ satisfying $\xi_0 = \eta$, where η satisfies $v_n(\eta) = \max v_n(x)$ and let $(x_i)_{i\in\mathbb{Z}}$ be the minimal configuration of $\bar{h}_n(x_i, x_{i+1}) = h_0(x_i, x_{i+1}) + u_n(x_{i+1})$ with rotation symbol 0^+ satisfying $x_0 \in [0,1]\backslash\Lambda_n$, then

$$\begin{aligned}
&\sum_{i\in\mathbb{Z}}(h_n(\xi_i,\xi_{i+1}) - h_n(\xi_i^-,\xi_{i+1}^-))\\
&\qquad\geq v_n(\eta) + \sum_{i\in\mathbb{Z}}\bar{h}_n(\xi_i,\xi_{i+1}) - \sum_{i\in\mathbb{Z}}h_n(\xi_i^-,\xi_{i+1}^-),\\
&\qquad\geq v_n(\eta) + \sum_{i\in\mathbb{Z}}\bar{h}_n(x_i,x_{i+1}) - \sum_{i\in\mathbb{Z}}h_n(x_i,x_{i+1}),\\
&\qquad= v_n(\eta) - \sum_{i\in\mathbb{Z}}v_n(x_{i+1}).
\end{aligned}$$

Therefore, we have shown:

$$P^{h_n}_{0^+}(\eta) = \min_{\xi_0=\eta}\sum_{i\in\mathbb{Z}}(h_n(\xi_i,\xi_{i+1}) - h_n(\xi_i^-,\xi_{i+1}^-)) \geq v_n(\eta) - \sum_{i\in\mathbb{Z}}v_n(x_{i+1}). \tag{2.1.44}$$

By (2.1.42), we have

$$v_n(\eta) \geq e^{-2N}n^{-a}. \tag{2.1.45}$$

It follows that

$$\sum_{i\in\mathbb{Z}}v_n(x_{i+1}) \leq \sigma^2e^{-2N}\sum_{i\in\mathbb{Z}}u_n(x_{i+1}) \leq \sigma^2e^{-2N}\sum_{i\in\mathbb{Z}}\frac{1}{4}(x_{i+1}-x_{i-1})^2 \leq \sigma^2e^{-2N}. \tag{2.1.46}$$

Hence,

$$P^{h_n}_{0^+}(\eta) \geq e^{-2N}(n^{-a} - \sigma^2), \tag{2.1.47}$$

we choose then σ (consequently N) in such a way that

$$\frac{1}{4}n^{-a} - \sigma^2 \geq 0,$$

which implies

$$\sigma \leq \frac{1}{2} n^{-\frac{a}{2}}.$$

By (3.4.4), if follows that

$$N \geq C n^{\frac{a}{2}+\frac{a}{2k}}. \tag{2.1.48}$$

Therefore,

$$\max_{N} P_{0^+}^{h_n}(\eta) \geq n^{-a} \exp\left(-C n^{\frac{a}{2}+\frac{a}{2k}}\right). \tag{2.1.49}$$

Second, following a similar argument as [W1], we have

$$|P_{\omega}^{h_n}(\xi) - P_{0^+}^{h_n}(\xi)| \leq C \exp\left(-2 n^{\frac{a}{2}+\frac{\delta}{2}}\right). \tag{2.1.50}$$

where $\xi \in \Lambda_n$ and δ is a small positive constant independent of n. Here Λ_n is as the same as the notation in (2.1.34).

Based on the preparations above, it is easy to prove Lemma 2.12. We assume that there exists an invariant circle with rotation number $0 < \omega < n^{-a-\delta}$ for h_n, then $P_{\omega}^{h_n}(\xi) \equiv 0$ for every $\xi \in \mathbb{R}$. By (2.1.50), we have

$$|P_{0^+}^{h_n}(\xi)| \leq C \exp\left(-2 n^{\frac{a}{2}+\frac{\delta}{2}}\right), \quad \text{for} \quad \xi \in \Lambda_n. \tag{2.1.51}$$

On the other hand, (2.1.49) implies that there exists a point $\eta \in \Lambda_n$ such that

$$P_{0^+}^{h_n}(\eta) \geq n^{-a} \exp\left(-C n^{\frac{a}{2}+\frac{a}{2k}}\right).$$

Hence, we have

$$n^{-a} \exp\left(-C n^{\frac{a}{2}+\frac{a}{2k}}\right) \leq C \exp\left(-2 n^{\frac{a}{2}+\frac{\delta}{2}}\right). \tag{2.1.52}$$

To achieve the contradiction, it suffices to take

$$k > \frac{3a}{2\delta},$$

which implies

$$\frac{a}{2k} < \frac{\delta}{3} < \frac{\delta}{2}.$$

Hence, for n large enough

$$n^{-a} \exp\left(-C n^{\frac{a}{2}+\frac{a}{2k}}\right) \geq C \exp\left(-2 n^{\frac{a}{2}+\frac{\delta}{2}}\right), \tag{2.1.53}$$

which contradicts (2.1.52). Therefore, there exists no invariant circle with rotation number $0 < \omega < n^{-a-\delta}$.

For $-n^{-a-\delta} < \omega < 0$, by comparing $P_{\omega}^{h_n}(\xi)$ with $P_{0^-}^{h_n}(\xi)$, the proof is similar. We omit the details. This completes the proof of Lemma 2.12. □

By Lemma 2.2, the case with a given irrational rotation number can be easily reduced to the one with a small enough rotation number. For the sake of simplicity of notations, we denote Q_{q_n} by Q_n and the same to u_{q_n}, v_{q_n} and h_{q_n}. Let

$$Q_n(x) = {q_n}^{-2}(u_n(q_n x) + v_n(q_n x)),$$

where $(q_n)_{n\in\mathbb{N}}$ is a sequence satisfying Dirichlet approximation

$$|q_n\omega - p_n| < \frac{1}{q_n}, \tag{2.1.54}$$

where $p_n \in \mathbb{Z}$ and $q_n \in \mathbb{N}$. Since $\omega \in \mathbb{R}\backslash\mathbb{Q}$, we say $q_n \to \infty$ as $n \to \infty$. Let $\tilde{h}_n(x, x') = h_0(x, x') + Q_n(x')$, we prove Theorem 2.11 for $(\tilde{h}_n)_{n\in\mathbb{N}}$ as follow:

Proof Based on Lemma 2.12 and Dirichlet approximation (2.1.54), it suffices to take

$$\frac{1}{q_n} \leq \frac{1}{{q_n}^{a+\delta}},$$

which implies

$$a \leq 1 - \delta. \tag{2.1.55}$$

From the constructions of u_n and v_n, it follows that

$$\begin{aligned}
&||\tilde{h}_n(x, x') - h_0(x, x')||_{C^r}\\
&\quad = ||Q_n(x')||_{C^r},\\
&\quad \leq {q_n}^{-2}(||u_n(q_n x')||_{C^r} + ||v_n(q_n x')||_{C^r}),\\
&\quad \leq {q_n}^{-2}({q_n}^{-a}(2\pi)^r {q_n}^r + C_1 {q_n}^{-a} {q_n}^r),\\
&\quad \leq C_2 {q_n}^{r-a-2},
\end{aligned}$$

where C_1, C_2 are positive constants only depending on r.

To complete the proof, it is enough to make $r - a - 2 < 0$, which together with (2.1.55) implies

$$r < a + 2 \leq 3 - \delta.$$

This completes the proof of Theorem 2.11. □

2.2. Case with $d \geq 3$ degrees of freedom

2.2.1. Preliminaries

In $\mathrm{T}^*\mathbb{T}^d$, a submanifold $\mathcal{T}^d$ is called Lagrangian torus if it is diffeomorphic to the torus $\mathbb{T}^d$ and the symplectic form vanishes on it. For positive definite Hamiltonian systems, if a Lagrangian torus is invariant under the Hamiltonian flow, it is then the graph over $\mathbb{T}^d$ (see [BP]). An example of Lagrangian torus is the KAM torus.

Definition 2.13 *$\bar{\mathcal{T}}^d$ is called d dimensional KAM torus if*

- *$\bar{\mathcal{T}}^d$ is a Lipschitz graph over $\mathbb{T}^d$;*
- *$\bar{\mathcal{T}}^d$ is invariant under the Hamiltonian flow Φ_t^H generated by the Hamiltonian function H;*
- *there exists a diffeomorphism $\phi:\ \mathbb{T}^d \to \bar{\mathcal{T}}^d$ such that $\phi^{-1} \circ \Phi_H^t \circ \phi = R_\omega^t$ for any $t \in \mathbb{R}$, where $R_\omega^t:\ x \to x + \omega t$ and ω is called the rotation vector of $\bar{\mathcal{T}}^d$.*

For positive definite Hamiltonian systems, each KAM torus $\bar{\mathcal{T}}^d$ supports a minimal measure μ. The rotation number ρ of μ is well defined and $\rho(\mu) = \omega$. The rotation vector of the Lagrangian torus $\mathcal{T}^d$ is not well defined. If $\mathcal{T}^d$ supports several invariant measures with different rotation vectors. In this paper, we are only concerned with Lagrangian tori with well defined rotation vectors.

Definition 2.14 *$\mathcal{T}^d$ is called d dimensional Lagrangian torus with the rotation vector ω if*

- *$\mathcal{T}^d$ is a Lagrangian submanifold;*
- *$\mathcal{T}^d$ is invariant for the Hamiltonian flow Φ_H^t generated by H.*
- *each orbit on $\mathcal{T}^d$ has the same rotation vector.*

In [H4], it is proved that each orbit on $\mathcal{T}^d$ is an action minimizing curve.

An arithmetic approximation of the rotation vector is found in [Ch]. For any given vector $\omega \in \mathbb{R}^d$ with $d \geq 2$, there is a sequence of integer vectors $k_n \in \mathbb{Z}^d$ with $|k_n| \to \infty$ such that

$$|\langle \omega, k_n \rangle| < \frac{C}{|k_n|^{d-1}}, \tag{2.2.1}$$

where C is a constant independent of n,

$$|k| = \left(\sum_{j=1}^{d} k_i^2 \right)^{\frac{1}{2}}, \quad \text{for } k = (k_1, k_2, \ldots, k_d).$$

A rotation vector $\omega \in \mathbb{R}^d$ is called resonant if there exists $k \in \mathbb{Z}^d$ such that $\langle \omega, k \rangle = 0$. Otherwise, it is non-resonant. Obviously, a Lagrangian torus with the resonant rotation vector can be destructed by analytic perturbation arbitrarily close to zero. Hence, it is sufficient to consider the Lagrangian torus with the non resonant rotation vector. In that case, one can assume that the Lagrangian torus $\mathcal{T}^d$ supports a uniquely ergodic minimizing measure. Moreover, by [Ma3], $\mathcal{T}^d$ is a Lipschitz graph over the underlying manifold $\mathbb{T}^d$.

2.2.2. C^∞ case

We will prove the following theorem:

Theorem 2.15 *Given an integrable Hamiltonian H_0 with d ($d \geq 3$) degrees of freedom, a rotation vector ω and a small positive constant δ, there exists a sequence of C^∞ Hamiltonians $\{H_n\}_{n\in\mathbb{N}}$ such that $H_n \to H_0$ in $C^{2d-\delta}$ topology and the Hamiltonian flow generated by H_n does not admit the Lagrangian torus with the rotation vector ω.*

This theorem implies that the rigidity of the Lagrangian torus is as the same as the KAM torus. Roughly speaking, the maximum of r is closely related to the arithmetic property of the rotation vector ω. If ω is a Diophantine vector, then r is at most $2d - \delta$. If ω is a Liouville vector, then r can be arbitrarily large. If ω can be approximated exponentially by rational vectors, then the Lagrangian torus with the rotation vector ω can be destructed by an arbitrarily small perturbation in C^ω (analytic) topology (see [Be]).

2.2.2.1. Destruction of Lagrangian torus with a special rotation vector

The Hamiltonian function we consider here is nearly integrable

$$H_n(q,p) = H_0(p) - P_n(q), \tag{2.2.2}$$

where $(q,p) \in \mathbb{T}^d \times \mathbb{R}^d$. Without loss of generality, we assume

$$H_0(p) = \frac{1}{2}|p|^2,$$

for which (2.2.2) is a typical mechanical system.

Since H_n is strictly convex with respect to p, by the Legendre transformation, the Lagrangian function corresponding to H_n is

$$L_n(q,\dot{q}) = \frac{1}{2}|\dot{q}|^2 + P_n(q), \tag{2.2.3}$$

where $\dot{q} = \frac{\partial H_0}{\partial p}$.

Let

$$P_n(q) = \frac{1}{n^a}(1 - \cos q_1) + v_n(q_1, q_2),$$

where a is a positive constant independent of n. For the rotation vector $\omega = (\omega_1, \dots, \omega_d)$, $v_n(q_1,q_2)$ is constructed as follow

$$\begin{cases} v_n \text{ is } 2\pi\text{-periodic}, \\ \operatorname{supp} v_n \cap \{[0,2\pi]\times[-\pi,\pi]\} = B_{R_n}(q^*), \\ \max_{(q_1,q_2)\in[0,2\pi]\times[-\pi,\pi]} v_n = v_n(q_0) = |\omega_1|^s, \\ ||v_n||_{C^r} \sim |\omega_1|^{s'}, \end{cases} \tag{2.2.4}$$

where $R_n = \frac{|\omega_1|}{n^2}$, $q^* = (\pi, 0)$ and we require $s' > 3$, it can be satisfied if $s > r + 3$.

For (2.2.2), we have the following lemma.

Lemma 2.16 *For n large enough, the Hamiltonian flow generated by $H_n(q,p)$ does not admit any Lagrangian torus with rotation vector $\omega = (\omega_1, \dots, \omega_d)$ satisfying*

$$|\omega_1| < n^{-\frac{a}{2}-\epsilon},$$

where $\epsilon > 0$ is independent of n.

Lemma 2.24 will be proved with variational method. First of all, we put it into the Lagrangian formalism. Let $\sigma_n = n^{-a}$. The Lagrangian function corresponding to (2.2.3) is

$$L_n\left(q_1, Q, \dot{q}_1, \dot{Q}\right) = \frac{1}{2}|\dot{Q}|^2 + \frac{1}{2}|\dot{q}_1|^2 + \sigma_n(1 - \cos(q_1)) + v_n(q_1, q_2), \tag{2.2.5}$$

where $Q = (q_2, \dots, q_d)$. $L_n(q_1, Q, \dot{q}_1, \dot{Q})$ can be considered as a perturbation coupling of a rotator with $d-1$ degrees of freedom and a perturbation with the Lagrangian function

$$A_n(q_1, \dot{q}_1) = \frac{1}{2}|\dot{q}_1|^2 + \sigma_n(1 - \cos(q_1)), \tag{2.2.6}$$

which corresponds to the Hamiltonian via Legendre transformation

$$h_n(q_1, p_1) = \frac{1}{2}|p_1|^2 - \sigma_n(1 - \cos q_1). \tag{2.2.7}$$

2.2.2.1.1. The action of the simple pendulum

Each solution of the Lagrangian equation determined by A_n, denoted by $q_1(t)$, determines an orbit $(q_1(t), p_1(t))$ of the Hamiltonian flow generated by h_n. Each orbit stays in certain energy level set $(q_1, p_1) \in h_n^{-1}(e)$. Under the boundary condition that $t_0 = 0,\ t_1 = \pi$ (or $t_1 = \pi, t_2 = 2\pi$), there is a unique correspondence between $t_1 - t_0$ and the energy, denoted by $e(t_1 - t_0)$, such that the determined orbit stays in the energy level set $h_n^{-1}(e(t_1 - t_0))$. More precisely, we have the following lemma.

Lemma 2.17 *Let $\bar{q}_1$ be the solution of A_n on $(t_0, \bar{t}_1)$ satisfying the boundary conditions*

$$\begin{cases} \bar{q}_1(t_0) = 0, \\ \bar{q}_1(\bar{t}_1) = \pi, \end{cases}$$

$e(\bar{t}_1 - t_0)$ be the energy of $\bar{q}_1$, i.e. $(\bar{q}_1, \bar{p}_1) \in h_n^{-1}(e(\bar{t}_1 - t_0))$ and ω_1 be the average speed of $\bar{q}_1$ on $(t_0, \bar{t}_1)$, then

$$e(\bar{t}_1 - t_0) \sim \sigma_n \exp\left(-\frac{C\sqrt{\sigma_n}}{|\omega_1|}\right), \tag{2.2.8}$$

where $f \sim g$ means that $\frac{1}{C}g < f < Cg$ holds for some constant $C > 0$, $\sigma_n = n^{-a}$.

Proof By the definition, we have

$$\frac{1}{2}|\dot{\bar{q}}_1|^2 - \sigma_n(1 - \cos(\bar{q}_1)) = e(\bar{t}_1 - t_0),$$

hence

$$|\dot{\bar{q}}_1| = \sqrt{2(e(\bar{t}_1 - t_0) + \sigma_n(1 - \cos(\bar{q}_1)))}.$$

Since the average speed of $\bar{q}_1$ is ω_1, by a direct calculation, we have

$$\frac{\pi}{|\omega_1|} = \int_{t_0}^{\bar{t}_1} dt = \int_0^{\pi} \frac{d\bar{q}_1}{\sqrt{2(e(\bar{t}_1 - t_0) + \sigma_n(1 - \cos(\bar{q}_1)))}} \sim \frac{1}{\sqrt{\sigma_n}} \ln\left(\frac{\sigma_n}{e(\bar{t}_1 - t_0)}\right),$$

moreover,

$$e(\bar{t}_1 - t_0) \sim \sigma_n \exp\left(-\frac{C\sqrt{\sigma_n}}{|\omega_1|}\right),$$

which complete the proof of Lemma 2.17. □

It is easy to see that Lemma 2.17 also holds for

$$\begin{cases} \bar{q}_1(\bar{t}_1) = \pi, \\ \bar{q}_1(t_2) = 2\pi. \end{cases}$$

The following lemma implies that the actions along orbits in the neighborhood of the separatix of the pendulum does not change too much with respect to a small change in speed (time).

Lemma 2.18 *Let $\bar{t}_1, \tilde{t}_1 \in [t_0, t_2]$. Let $\bar{q}_1(t)$ be a solution of A_n on $(t_0, \bar{t}_1)$ and $(\bar{t}_1, t_2)$ with boundary conditions respectively*

$$\begin{cases} \bar{q}_1(t_0) = 0, \\ \bar{q}_1(\bar{t}_1) = \pi, \end{cases} \quad \begin{cases} \bar{q}_1(\bar{t}_1) = \pi, \\ \bar{q}_1(t_2) = 2\pi, \end{cases}$$

and let $\tilde{q}_1(t)$ *be a solution of* A_n *on* $(t_0, \tilde{t}_1)$ *and* $(\tilde{t}_1, t_2)$ *with boundary conditions respectively*

$$\begin{cases} \tilde{q}_1(t_0) = 0, \\ \tilde{q}_1(\tilde{t}_1) = \pi, \end{cases} \quad \begin{cases} \tilde{q}_1(\tilde{t}_1) = \pi, \\ \tilde{q}_1(t_2) = 2\pi. \end{cases}$$

Let $\bar{\omega}_1'$ *and* $\bar{\omega}_1''$ *be the average speed of* $\bar{q}_1$ *on* $(t_0, \bar{t}_1)$ *and* $(\bar{t}_1, t_2)$ *respectively. Let* $\tilde{\omega}_1'$ *and* $\tilde{\omega}_1''$ *be the average speed of* $\tilde{q}_1$ *on* $(t_0, \tilde{t}_1)$ *and* $(\tilde{t}_1, t_2)$ *respectively. We set*

$$|\omega_1| = \max\left\{|\bar{\omega}_1'|, |\bar{\omega}_1''|, |\tilde{\omega}_1'|, |\tilde{\omega}_1''|\right\},$$

then

$$\left|\int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt - \int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt\right| \leq C_1 |\bar{t}_1 - \tilde{t}_1| \sigma_n \exp\left(-\frac{C_2\sqrt{\sigma_n}}{|\omega_1|}\right). \tag{2.2.9}$$

Proof The proof follows the similar idea of Lemma 4 in [Be]. Let $q_1(t)$ be a solution of A_n on (t_0, t_1) and (t_1, t_2) with boundary conditions respectively

$$\begin{cases} q_1(t_0) = 0, \\ q_1(t_1) = \pi, \end{cases} \quad \begin{cases} q_1(t_1) = \pi, \\ q_1(t_2) = 2\pi. \end{cases}$$

We consider the function

$$\begin{aligned} L(t_1) = & \int_{t_0}^{t_1} A_n(q_1, \dot{q}_1)dt + \int_{t_1}^{t_2} A_n(q_1, \dot{q}_1)dt, \\ = & \int_0^{\pi} \sqrt{2(e(t_1 - t_0) + V(q_1))}dq_1 - e(t_1 - t_0)(t_1 - t_0) \\ & + \int_{\pi}^{2\pi} \sqrt{2(e(t_2 - t_1) + V(q_1))}dq_1 - e(t_2 - t_1)(t_2 - t_1), \end{aligned}$$

where

$$V(q_1) = \sigma_n(1 - \cos(q_1)),$$

and $e(\Delta t)$ denotes the energy of the orbit of the pendulum moving half a turn in time Δt. The quantity $e(\Delta t)$ is differentiable with respect to Δt, then

$$\begin{aligned} \frac{dL(t_1)}{dt_1} = & \int_0^{\pi} \frac{\dot{e}(t_1 - t_0)}{\sqrt{2(e(t_1 - t_0) + V(q_1))}}dq_1 - \dot{e}(t_1 - t_0)(t_1 - t_0) \\ & - e(t_1 - t_0) - \int_{\pi}^{2\pi} \frac{\dot{e}(t_2 - t_1)}{\sqrt{2(e(t_2 - t_1) + V(q_1))}}dq_1 \\ & + \dot{e}(t_2 - t_1)(t_2 - t_1) + e(t_2 - t_1), \\ = & \int_{t_0}^{t_1} \dot{e}(t_1 - t_0)dt - \dot{e}(t_1 - t_0)(t_1 - t_0) - e(t_1 - t_0) \\ & - \int_{t_1}^{t_2} \dot{e}(t_2 - t_1)dt + \dot{e}(t_2 - t_1)(t_2 - t_1) + e(t_2 - t_1), \\ = & e(t_2 - t_1) - e(t_1 - t_0). \end{aligned} \tag{2.2.10}$$

Thus, we have

$$\left|\frac{dL(t_1)}{dt_1}\right| \leq |e(t_2 - t_1)| + |e(t_1 - t_0)|.$$

Integrate from $\bar{t}_1$ to $\tilde{t}_1$ and from (2.2.8), it follows that

$$\left|\int_{t_0}^{t_2} A_n(\bar{q}_1,\dot{\bar{q}}_1)dt - \int_{t_0}^{t_2} A_n(\tilde{q}_1,\dot{\tilde{q}}_1)dt\right| \le C_1|\bar{t}_1-\tilde{t}_1|\sigma_n \exp\left(-\frac{C_2\sqrt{\sigma_n}}{|\omega_1|}\right),$$

which completes the proof of Lemma 2.18. □

2.2.2.1.2. The velocity of the action minimizing orbit

Once the function $q_1(t)$ is fixed, the function $Q(t)$ is the solution of the Euler-Lagrange equation with the non autonomous Lagrangian

$$\frac{1}{2}|\dot{Q}(t)|^2 + v_n\left(q_1(t),q_2(t)\right), \tag{2.2.11}$$

where $Q(t)=(q_2(t),\dots,q_d(t))$.

Lemma 2.19 *Let $(q_1(t),Q(t))$ be the orbit of L_n with rotation vector ω, then for any $t',t''\in\mathbb{R}$ and $t\in[t',t'']$ we have*

$$\left|\dot{Q}(t)-\frac{Q(t'')-Q(t')}{t''-t'}\right| \le C|\omega_1|^2. \tag{2.2.12}$$

Proof By the Euler-Lagrange equation, we have

$$\ddot{q}_i(t)=0, \quad \text{for} \quad i=3,\dots,d,$$

hence $\dot{q}_i(t)=\text{const.}$, (2.2.12) is verified obviously for $q_i(t)$, $i=3,\dots,d$. We just need to consider $q_2(t)$. Let $q_1(t_0)=0$ and $q_1(t_2)=2\pi$. It suffices to prove that for $t\in[t',t'']\subset[t_0,t_2]$

$$\left|\dot{q}_2(t)-\frac{q_2(t'')-q_2(t')}{t''-t'}\right| \le C|\omega_1|^2. \tag{2.2.13}$$

From the Euler-Lagrange equation,

$$\ddot{q}_2(t)=\frac{\partial v_n}{\partial q_2}(q_1(t),q_2(t)),$$

together with $||v_n||_{C^r}\sim|\omega_1|^{s'}$, we obtain

$$\ddot{q}_2(t)\le C_1|\omega_1|^{s'}.$$

Integrate the two sides of the inequality above from t' to t'', we have

$$|\dot{q}_2(t'')-\dot{q}_2(t')|\le C_2|\omega_1|^{s'}|t''-t'|.$$

It follows from (2.2.1) that $|t''-t'|\le C_3|\omega_1|^{-1}$. Hence

$$|\dot{q}_2(t'')-\dot{q}_2(t')|\le C_4|\omega_1|^{s'-1}.$$

Since $s'>3$, we have

$$|\dot{q}_2(t'')-\dot{q}_2(t')|\le C_4|\omega_1|^2.$$

This completes the proof. □

2.2.2.1.3. Proof of Lemma 2.24

Based on the minimal property of the orbits on an invariant Lagrangian torus and its graph property, passing through each $x \in \mathbb{T}^d$, there is a unique minimal curve $q(t)$ with rotation vector ω if the Hamiltonian flow generated by H_n admits a Lagrangian torus with rotation vector ω. Hence, it is sufficient to prove the existence of some point in $\mathbb{T}^d$ where no minimal curve passes through.

Indeed, any minimal curve does not pass through the subspace $(\pi, 0) \times \mathbb{T}^{d-2}$. It implies Lemma 2.24. Let us assume the contrary, namely, there exists $\bar{t}_1$ such that

$$q_1(\bar{t}_1) = \pi, \quad q_2(\bar{t}_1) = 0,$$

where $q(t) = (q_1, q_2, \dots, q_d)(t)$ is a minimal curve in the universal covering space $\mathbb{R}^d$. Because of $\omega_1 \neq 0$, there exist t_0 and t_2 such that

$$q_1(t_0) = 0, \quad q_1(t_2) = 2\pi.$$

Obviously, $t_0 < \bar{t}_1 < t_2$ and

$$t_2 - t_0 \sim \frac{1}{|\omega_1|}.$$

Let $\tilde{t}_1$ be the last time before $\bar{t}_1$ or the first time after $\bar{t}_1$ such that

$$|q_2(\tilde{t}_1) - q_2(\bar{t}_1)| = \pi.$$

It is easy to see that,

$$|\tilde{t}_1 - \bar{t}_1| \sim \frac{1}{|\omega_2|}.$$

Since $|\omega_2| \sim 1$, then

$$|\tilde{t}_1 - \bar{t}_1| \leq C_0.$$

Without loss of generality, one can assume $\omega_1 > 0$ and $\omega_2 > 0$. Consider a solution $\tilde{q}_1$ of A_n on $(t_0, \tilde{t}_1)$ and on $(\tilde{t}_1, t_2)$ with boundary conditions respectively

$$\begin{cases} \tilde{q}_1(t_0) = q_1(t_0) = 0, \\ \tilde{q}_1(\tilde{t}_1) = q_1(\bar{t}_1) = \pi, \end{cases} \qquad \begin{cases} \tilde{q}_1(\tilde{t}_1) = q_1(\bar{t}_1) = \pi, \\ \tilde{q}_1(t_2) = q_1(t_2) = 2\pi. \end{cases}$$

Since q is assumed to be a minimal curve, we have

$$\int_{t_0}^{t_2} L_n(\tilde{q}_1, Q, \dot{\tilde{q}}_1, \dot{Q})dt - \int_{t_0}^{t_2} L_n(q_1, Q, \dot{q}_1, \dot{Q})dt \geq 0. \tag{2.2.14}$$

See Fig.1, where $x_1 = (q_1(\bar{t}_1), q_2(\bar{t}_1)) = (\pi, 0)$, $x_0 = (q_1(t_0), q_2(t_0)) = (0, q_2(t_0))$, $x_2 = (q_1(t_2), q_2(t_2)) = (2\pi, q_2(t_2))$, $\tilde{x}_1' = (\pi, -\pi)$ and $\tilde{x}_1'' = (\pi, \pi)$.

$(\tilde{q}_1(t), q_2(t))$ passes through the point $\tilde{x}_1'$ or $\tilde{x}_1''$. Thus, by the construction of L_n, we obtain from (2.2.14) that

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt \geq \int_{t_0}^{t_2} v_n(q_1, q_2)dt - \int_{t_0}^{t_2} v_n(\tilde{q}_1, q_2)dt. \tag{2.2.15}$$

By the definition of v_n as (2.2.4), we find

$$(\tilde{q}_1(t), q_2(t)) \cap \operatorname{supp} v_n = \emptyset, \quad \text{for} \quad t \in (t_0, t_2).$$



Figure 1: The projections of the curves $(q_1(t), Q(t))$ and $(\tilde{q}_1(t), Q(t))$ on $[0, 2\pi] \times \mathbb{R}$

In fact, if there would exist $\hat{t}$ such that $(\tilde{q}_1(\hat{t}), q_2(\hat{t})) \in \operatorname{supp} v_n$, without loss of generality, one can assume $\hat{t} > \tilde{t}_1$. By Lemma 2.19, for any $t \in [\tilde{t}_1, \hat{t}]$,

$$\dot{q}_2(t) \le C_1,$$

hence,

$$\hat{t} - \tilde{t}_1 \ge C_2,$$

where C_1, C_2 are constants independent of n. Consequently

$$|\tilde{q}_1(\hat{t}) - \tilde{q}_1(\tilde{t}_1)| \ge C_3|\omega_1| > R_n,$$

where R_n is the radius of the support of v_n. It is impossible.

Hence, we have

$$\int_{t_0}^{t_2} v_n(q_1, q_2)dt - \int_{t_0}^{t_2} v_n(\tilde{q}_1, q_2)dt = \int_{t_0}^{t_2} v_n(q_1, q_2)dt.$$

By the construction of v_n and the minimality of (q_1, Q), a simple calculation shows

$$\int_{t_0}^{t_2} v_n(q_1, q_2)dt \ge |\omega_1|^{\lambda}, \tag{2.2.16}$$

where λ is a positive constant. Consequently, if follows from (2.2.15) that

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt \ge |\omega_1|^{\lambda}. \tag{2.2.17}$$

On the other hand, consider a solution $\bar{q}_1$ of A_n on $(t_0, \bar{t}_1)$ and on $(\bar{t}_1, t_2)$ with boundary conditions respectively

$$\begin{cases} \bar{q}_1(t_0) = q_1(t_0) = 0, \\ \bar{q}_1(\bar{t}_1) = q_1(\bar{t}_1) = \pi, \end{cases} \qquad \begin{cases} \bar{q}_1(\bar{t}_1) = q_1(\bar{t}_1) = \pi, \\ \bar{q}_1(t_2) = q_1(t_2) = 2\pi. \end{cases}$$

Along both of which the action of A_n achieves its minimum. Thus, we have

$$\int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt \ge \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt.$$

We compare the action $\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt$ with the action $\int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt$ in the alternative cases, which is based on the choices of $\tilde{t}_1$. See Fig.2, where $\bar{t} = \frac{t_0+t_2}{2}$.



Case 1 Case 2a Case 2b

Figure 2: The choices of $\tilde{t}_1$

Case 1: $|\bar{t}_1 - \bar{t}| \leq C_0$.

In this case, the average speed of $\bar{q}_1$ on $(t_0, \bar{t}_1)$ and $(\bar{t}_1, t_2)$ have the same quantity order as $|\omega_1|$. By $|\tilde{t}_1 - \bar{t}_1| \leq C_0$, we have $|\tilde{t}_1 - \bar{t}| \leq 2C_0$. Hence the average speed of $\tilde{q}_1$ on $(t_0, \tilde{t}_1)$ and $(\tilde{t}_1, t_2)$ have also the same quantity order as $|\omega_1|$. Thus, Lemma 2.18 implies

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt \leq C_4\sigma_n \exp\left(-\frac{C_5\sqrt{\sigma_n}}{|\omega_1|}\right).$$

Case 2: $|\bar{t}_1 - \bar{t}| > C_0$.

In this case, we take $\tilde{t}_1$ such that $|\tilde{t}_1 - \bar{t}| \leq |\bar{t}_1 - \bar{t}|$, which can be achieved by the suitable choice of the position of $\tilde{q}_1(\tilde{t}_1)$. More precisely,

- if $\bar{t}_1 > \bar{t} + C_0$ (Case 2a in Fig.2), we choose $\tilde{t}_1$ as the last time before $\bar{t}_1$, i.e. $(\tilde{q}_1(\tilde{t}_1), q_2(\tilde{t}_1)) = \tilde{x}'_1$ in Fig.1;
- if $\bar{t}_1 < \bar{t} - C_0$ (Case 2b in Fig.2), we choose $\tilde{t}_1$ as the first time after $\bar{t}_1$, i.e. $(\tilde{q}_1(\tilde{t}_1), q_2(\tilde{t}_1)) = \tilde{x}''_1$ in Fig.1.

For both cases 2a and 2b, it follows from (2.2.10) that

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt \leq 0.$$

Hence, for any $\bar{t}_1 \in (t_0, t_2)$, we can find $\tilde{t}_1$ such that

$$\begin{aligned}\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt &\leq \int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt, \\ &\leq C_4\sigma_n \exp\left(-\frac{C_5\sqrt{\sigma_n}}{|\omega_1|}\right).\end{aligned}$$

Since

$$|\omega_1| \leq n^{-\frac{a}{2}-\epsilon}.$$

It is easy to see that for n large enough,

$$C_4\sigma_n \exp\left(-\frac{C_5\sqrt{\sigma_n}}{|\omega_1|}\right) \leq |\omega_1|^\lambda,$$

where $\sigma_n = n^{-a}$, which contradicts to (2.2.17) for large n. This completes the proof of Lemma 2.24. □

2.2.2.2. Destruction of Lagrangian torus with an arbitrary rotation vector

By (2.2.1), for every non resonant rotation vector $\omega = (\omega_1, \ldots, \omega_d)$ $(d \geq 2)$, there exists a sequence of integer vector $k_n \in \mathbb{Z}^d$ satisfying $|k_n| \to \infty$ as $n \to \infty$ such that

$$|\langle k_n, \omega\rangle| < \frac{C}{|k_n|^{d-1}}.$$

2.2.2.2.1. Transformation of coordinates

We choose a sequence of $k_n \in \mathbb{Z}^d$ satisfying (2.2.1) and an integer vector sequence k'_n such that $\langle k'_n, k_n\rangle = 0$. In addition, select $d-2$ integer vectors $l_{n3}, \ldots, l_{nd}$ such that $k_n, k'_n, l_{n3}, \ldots, l_{nd}$ are pairwise orthogonal. Let

$$K_n = (k_n, k'_n, l_{n3}, \ldots, l_{nd})^t. \tag{2.2.18}$$

We choose the transformation of the coordinates

$$q = K_n x.$$

Let p denotes the dual coordinate of q in the sense of Legendre transformation, i.e. $p = \frac{\partial L}{\partial \dot{q}}$, it follows that

$$y = K_n^t p,$$

where K_n^t denotes the transpose of K_n. We set

$$\Phi_n = \begin{pmatrix} K_n & \\ & K_n^{-t} \end{pmatrix},$$

then

$$\begin{pmatrix} q \\ p \end{pmatrix} = \Phi_n \begin{pmatrix} x \\ y \end{pmatrix}.$$

It is easy to verify that

$$\Phi_n^t J_0 \Phi_n = J_0,$$

where

$$J_0 = \begin{pmatrix} \mathbf{0} & \mathbf{1} \\ -\mathbf{1} & \mathbf{0} \end{pmatrix},$$

where $\mathbf{1}$ denotes a $d \times d$ unit matrix. Hence, Φ_n is a symplectic transformation in the phase space.

Lemma 2.20 *If the Hamiltonian flow generated by $\tilde{H}_n(x, y)$ admits a Lagrangian torus with rotation vector ω, then the Hamiltonian flow generated by $H_n(q, p)$ also admits a Lagrangian torus with rotation vector $K_n\omega$, where $(q, p)^t = \Phi_n(x, y)^t$.*

Proof Let $\tilde{\mathcal{T}}^d$ be the Lagrangian torus admitted by $\tilde{H}_n(x, y)$, a symplectic form Ω vanishes on $\mathrm{T}_x\tilde{\mathcal{T}}^d$ for every $x \in \tilde{\mathcal{T}}^d$. Since K_n consists of integer vectors, then $\mathcal{T}^d := K_n\tilde{\mathcal{T}}^d$ is still a torus. Φ_n is a symplectic transformation, hence $\mathcal{T}^d$ is a Lagrangian torus. From Definition 3.1, each orbit on $\tilde{\mathcal{T}}^d$ has the same rotation vector ω. Let $\tilde{\gamma}(t)$ be a lift of an orbit on $\tilde{\mathcal{T}}^d$, it follows that

$$\omega = \lim_{t \to \infty} \frac{\tilde{\gamma}(t) - \tilde{\gamma}(-t)}{2t}.$$

For $\gamma(t) = K_n\tilde{\gamma}(t)$, we have

$$\begin{aligned}\lim_{t\to\infty}\frac{\gamma(t)-\gamma(-t)}{2t} &= \lim_{t\to\infty}\frac{K_n\tilde{\gamma}(t)-K_n\tilde{\gamma}(-t)}{2t};\\ &= K_n\lim_{t\to\infty}\frac{\tilde{\gamma}(t)-\tilde{\gamma}(-t)}{2t};\\ &= K_n\omega.\end{aligned}$$

This completes the proof. □

Remark 2.21 *For $\tilde{\mathcal{T}}^d$ and $\mathcal{T}^d$ in the proof of Lemma 2.20, we have*

$$Vol(\mathcal{T}^d) = |\det K_n|\, Vol(\tilde{\mathcal{T}}^d),$$

where Vol(·) denotes the volume of (·).

2.2.2.2.2. Proof of Theorem 2.15

We construct $\tilde{H}_n(x,y)$ as follow:

$$\tilde{H}_n(x,y) = \frac{1}{2}|y|^2 - \tilde{P}_n(x), \tag{2.2.19}$$

where

$$\tilde{P}_n(x) = \frac{1}{|k_n|^{a+2}}(1-\cos\langle k_n,x\rangle) + \frac{1}{|k_n|^2}v_n\left(\langle k_n,x\rangle, \langle k_n',x\rangle\right),$$

where k_n' is the second row of K_n and v_n is defined by (2.2.4). Let $q = K_n x$. In particular, we have

$$\begin{cases} q_1 = \langle k_n, x\rangle, \\ q_2 = \langle k_n', x\rangle. \end{cases} \tag{2.2.20}$$

By the transformation of coordinates and the Legendre transformation, the Lagrangian function corresponding to (2.2.19) is

$$\begin{aligned} L_n\left(q_1,Q,\dot{q}_1,\dot{Q}\right) =& \frac{1}{2}\sum_{i=3}^{d}\frac{|\dot{q}_i|^2}{|l_{ni}|^2} + \frac{|\dot{q}_2|^2}{2|k_n'|^2} \\ &+ \frac{1}{|k_n|^2}\left(\frac{1}{2}|\dot{q}_1|^2 + \frac{1}{|k_n|^a}(1-\cos(q_1)) + v_n(q_1,q_2)\right), \end{aligned} \tag{2.2.21}$$

where $Q = (q_2,\dots,q_d)$.

For the Hamiltonian flow generated by (2.2.19), by Lemma 2.20, for the destruction of the Lagrangian torus $\tilde{\mathcal{T}}^d$ with rotation vector ω, it is sufficient to prove that the Euler-Lagrange flow generated by (2.2.21) admits no the Lagrangian torus $\mathcal{T}^d := K_n\tilde{\mathcal{T}}^d$ with rotation vector $K_n\omega$. Let $K_n\omega = (\omega_1,\omega_2,\dots,\omega_d)$.

It is easy to see that there exists an integer vector k_n' such that

$$\langle k_n, k_n'\rangle = 0 \quad \text{and} \quad |\langle k_n',\omega\rangle| \sim 1, \tag{2.2.22}$$

i.e. $\omega_2 \sim 1$. In fact, it suffices to consider $k_n' \in \mathbb{Z}^3$. Let $k_n' = (k_{n1}', k_{n2}', k_{n3}')$, then for $k_n' \in \mathbb{Z}^d$, one can take $k_n' = (k_{n1}', k_{n2}', k_{n3}', 0, \dots, 0)$ to verify (2.2.22).

Since ω is non-resonant, then $|k_n'| \to \infty$, for $n \to \infty$. Let θ be the angle between k_n' and ω, then

$$|\langle k_n', \omega\rangle| = |k_n'||\omega|| \cos\theta|,$$

where $|k_n'|$ is determined by (2.2.24) below. Let Π be the plane orthogonal with respect to k_n. Let $S_{R\alpha} \subset \Pi$ be the sector with cental point $(0, 0, 0)$, central angle α and radius R satisfying $\alpha = \frac{C_1}{|k_n'|}$ and the angle between ω and one of the radii is equal to $\frac{\pi}{2} - \frac{C_2}{|k_n'|}$, where $C_2 > C_1$. To achieve (2.2.22), it is sufficient to find an integer point $m = (m_1, m_2, m_3) \in \mathbb{Z}^3$ satisfying

$$|m| \sim |k_n'| \quad \text{and} \quad m \in S_{R\alpha}. \tag{2.2.23}$$

In deed, we take $k_n' = (m_1 - 0, m_2 - 0, m_3 - 0)$. Since

$$\cos\theta \sim \cos(\frac{\pi}{2} - \frac{1}{|k_n'|}) = \sin\frac{1}{|k_n'|} \sim \frac{1}{|k_n'|},$$

we have $|\langle k_n', \omega\rangle| \sim 1$. It is easy to see that there exists a suitable constant $r(|k_n'|)$ only depending on $|k_n'|$ such that the square of area $(r(|k_n'|))^2$ contains at least one integer point. We take

$$|k_n'| \sim (r(|k_n'|))^\kappa, \tag{2.2.24}$$

where $\kappa \gg 1$, then it can be concluded that the integer satisfying (2.2.34) does exist.

Replacing n by $|k_n|$ in the proof of Lemma 2.24, we have that the Euler-Lagrange flow generated by (2.2.21) does not admit any Lagrangian torus with rotation vector satisfying

$$|\omega_1| < |k_n|^{-\frac{a}{2}-\epsilon}.$$

From the construction of K_n, $|\omega_1| = |\langle k_n, \omega\rangle|$ which together with (2.2.1) implies

$$|\omega_1| < \frac{C}{|k_n|^{d-1}}.$$

Based on Lemma 2.20, it suffices to take

$$\frac{C}{|k_n|^{d-1}} \leq |k_n|^{-\frac{a}{2}-\epsilon},$$

which implies

$$a < 2d - 2 - 2\epsilon. \tag{2.2.25}$$

It follows from (2.2.4) and (2.2.19) that

$$\begin{aligned}
&||\tilde{H}_n(x, y) - H_0(y)||_{C^r} \\
&\quad = ||\tilde{P}_n(x)||_{C^r}, \\
&\quad = |k_n|^{-2} \left(|k_n|^{-a}||1 - \cos\langle k_n, x\rangle||_{C^r} + ||v_n(\langle k_n, x\rangle, \langle k_n', x\rangle)||_{C^r}\right), \\
&\quad \leq |k_n|^{-2} \left(C_1|k_n|^{-a+r} + C_2|k_n|^{-s'(d-1)+r}\right), \\
&\quad \leq C_3 \left(|k_n|^{r-a-2} + |k_n|^{r-3(d-1)-2}\right).
\end{aligned}$$

To complete the proof of Theorem 2.15, it is enough to make $r - a - 2 < 0$ and $r - 3(d - 1) - 2 < 0$, which together with (2.2.25) implies

$$r < 2d - 2\epsilon.$$

Taking $\delta = 3\epsilon$, this completes the proof of Theorem 2.15. □

2.2.3. C^ω case

We will prove the following theorem:

Theorem 2.22 *Given an integrable Hamiltonian H_0 with d ($d \geq 3$) degrees of freedom, a rotation vector ω and a small positive constant δ, there exists a sequence of C^ω Hamiltonians $\{H_n\}_{n\in\mathbb{N}}$ such that $H_n \to H_0$ in $C^{d+1-\delta}$ topology and the Hamiltonian flow generated by H_n admits no Lagrangian torus with the rotation vector ω.*

2.2.3.1. Construction of H_n

$P_n(x)$ is constructed as follow:

$$P_n(x) = \frac{1}{|k_n|^{d+1-\epsilon}}(1-\cos\langle k_n, x\rangle) + \mu_n \frac{1}{|k_n|^{d+1-\epsilon}}(1-\cos\langle k_n, x\rangle)\cos\langle k_n', x\rangle,$$

where k_n, k_n' are the first two rows of K_n defined as (2.2.18), ϵ is a given small positive constant and μ_n satisfies

$$\mu_n \sim \exp\left(-|k_n|^{\frac{d}{2}-\frac{1}{2}+\frac{\epsilon}{3}}\right). \tag{2.2.26}$$

A simple calculation implies that for $\delta = 3\epsilon$

$$||H_n(x,y) - H_0(y)||_{C^{d+1-\delta}} = ||P_n(x)||_{C^{d+1-\delta}} \to 0 \quad \text{as } n \to \infty.$$

From the transformation matrix of coordinates (2.2.18), let $\delta_n = \frac{1}{|k_n|^{d-1-\epsilon}}$, the Lagrangian function corresponding to (2.2.3) is

(2.2.27)
$$\begin{aligned} L_n\left(q_1, Q, \dot{q}_1, \dot{Q}\right) = {} & \frac{1}{2}\sum_{i=3}^{d}\frac{|\dot{q}_i|^2}{|l_{ni}|^2} + \frac{|\dot{q}_2|^2}{2|k_n'|^2} + \frac{1}{|k_n|^2}\left(\frac{1}{2}|\dot{q}_1|^2 + \delta_n(1-\cos(q_1))\right) \\ & + \frac{1}{|k_n|^2}\left(\mu_n\delta_n(1-\cos(q_1))\cos(q_2)\right), \end{aligned}$$

where $Q = (q_2, \ldots, q_d)$.

$L_n(q_1, Q, \dot{q}_1, \dot{Q})$ can be considered as a perturbation coupling of a rotator with $d-1$ degrees of freedom and a perturbation with the Lagrangian function

$$A_n(q_1, \dot{q}_1) = \frac{1}{2}|\dot{q}_1|^2 + \delta_n(1-\cos(q_1)), \tag{2.2.28}$$

which corresponds to the Hamiltonian via Legendre transformation

$$h_n(q_1, p_1) = \frac{1}{2}|p_1|^2 - \delta_n(1-\cos q_1). \tag{2.2.29}$$

We denote the coupling perturbation by

$$\tilde{P}_n(q_1, Q) = \mu_n\delta_n(1-\cos(q_1))\cos(q_2). \tag{2.2.30}$$

2.2.3.2. Melnikov function

In the following, we give some approximation lemmas on the actions of (2.2.28) and (2.2.30) along the minimal orbits of L_n by the calculation of Melnikov function.

For $t \in \mathbb{R}$, the separatrix $\hat{q}_1(t)$ of A_n as (2.2.28) satisfying $\dot{\hat{q}}_1 > 0$ and $\hat{q}_1(0) = \pi$ is

$$\begin{cases} \hat{q}_1(t) = 4\arctan\left(\exp(\sqrt{\delta_n}t)\right), \\ \dot{\hat{q}}_1(t) = \frac{2\sqrt{\delta_n}}{\cosh(\sqrt{\delta_n}t)}. \end{cases} \tag{2.2.31}$$

Let the separatrix of A_n takes value π at t_1 and $Q(t_1) = Q_1$, then for a given rotation vector ω, we define the Melnikov function as

$$M_n(\omega, Q_1, t_1) = \delta_n \int_{\mathbb{R}} (1 - \cos(\hat{q}_1(t - t_1))) \cos\left(\langle k_n', \omega(t - t_1)\rangle + q_2(t_1)\right) dt.$$

Namely, M_n is the integral of the coupling perturbation (2.2.30) along the separatrix of A_n. It can be explicitly calculated as follow

$$M_n(\omega, Q_1, t_1) = 2\pi \frac{\langle \omega, k_n'\rangle}{\sinh(\frac{\pi\langle\omega, k_n'\rangle}{2\sqrt{\delta_n}})} \cos(q_2(t_1)). \tag{2.2.32}$$

It is easy to see that M_n only depends on $\omega_2 = \langle \omega, k_n'\rangle$ and $q_2(t_1)$ on which M_n is 2π periodic. For the simplicity of notations, we denote $M_n(\omega, Q_1, t_1)$ by

$$M_n(\omega_2, Q_1, t_1) = 2\pi \frac{\omega_2}{\sinh(\frac{\pi\omega_2}{2\sqrt{\delta_n}})} \cos(q_2(t_1)). \tag{2.2.33}$$

Next, we work on the universal covering space of $\mathbb{T}^d$. By (2.2.33), a simple calculation implies the next lemma

Lemma 2.23 *If $\bar{q}_2(t') \,mod\, 2\pi = 0$ and $\tilde{q}_2(t'') \,mod\, 2\pi = \pi$. Let*

$$Q' = (\bar{q}_2(t'), q_3(t'), \dots, q_d(t')) \quad \text{and} \quad Q'' = \tilde{q}_2(t''), q_3(t''), \dots, q_d(t'')),$$

then for n large enough

$$M_n(\omega_2, Q', t') - M_n(\omega_2, Q'', t'') \geq \exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right), \tag{2.2.34}$$

where λ is a positive constant independent of n.

Proof By (2.2.22), we have

$$|\omega_2| \sim 1.$$

A simple calculation gives

$$M_n(\omega_2, Q', t') - M_n(\omega_2, Q'', t'') \geq \exp\left(-\frac{C_2|\omega_2|}{\sqrt{\delta_n}}\right) \geq \exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right),$$

where λ is a positive constant independent of n. This completes the proof of (2.2.34). □

2.2.3.3. An approximation lemma

We use M_n to approximate the action of the perturbation (2.2.30) along the minimal orbits of L_n.

Lemma 2.24 *Let* $(q_1(t), Q(t))$ *be the minimal orbit of* L_n *satisfying* $q_1(t_1) = \pi$ *with rotation vector* ω*, then*

(i) *there exist* $\tau > 0$ *and* $t_0,\ t_2$ *satisfying*

$$t_0 \leq t_1 - \frac{\tau}{\delta_n'} < t_1 + \frac{\tau}{\delta_n'} \leq t_2, \tag{2.2.35}$$

such that

$$q_1(t_0) = 0, \quad q_1(t_2) = 2\pi.$$

where

$$\delta_n' = \frac{1}{|k_n|^{d-1-\frac{\epsilon}{4}}};$$

(ii) *let*

$$\bar{\omega} = \left(\omega_1, \frac{Q(t_2) - Q(t_0)}{t_2 - t_0}\right), \tag{2.2.36}$$

then

$$\left|\delta_n \int_{t_0}^{t_2} (1 - \cos(q_1(t))) \cos(q_2(t)) dt - M_n(\bar{\omega}_2, Q(t_1), t_1)\right| \\ \leq C\sqrt{\delta_n} \exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right),$$

where C *is a positive constant independent of* n *and* λ *is the same as the one in (2.2.34).*

Proof The proof of Lemma 2.24 follows from the ideas of [Be]. We will prove (i) and (ii) respectively in the following. For the simplicity of notations, we will use $u \preceq v$ (resp. $u \succeq v$) to denote $u \leq Cv$ (resp. $u \geq Cv$) for some positive constant C.

2.2.3.3.1. Proof of (i)

We set $\tau = \frac{\pi}{C_0}$, where C_0 is the constant in (2.2.1). From (2.2.1), it follows that

$$\frac{2\tau}{\delta_n'} < \frac{2\pi}{|\omega_1|}.$$

Hence, we have either

$$t_0 \leq t_1 - \frac{\tau}{\delta_n'} \quad \text{or} \quad t_1 + \frac{\tau}{\delta_n'} \leq t_2.$$

Without loss of generality, we assume $t_1 + \frac{\tau}{\delta_n'} \leq t_2$ and prove $t_0 \leq t_1 - \frac{\tau}{\delta_n'}$ in the following.

Let t_{-1} and t_3 be the last time to the left of t_1 such that $q_1(t_{-1}) = -\pi$ and the first time to the right of t_1 such that $q_1(t_3) = 3\pi$ respectively. Consider the solution $\bar{q}$ of A_n on (t_{-1}, t_1) satisfying the boundary condition

$$\begin{cases} \bar{q}(t_{-1}) = q_1(t_{-1}) = -\pi, \\ \bar{q}(t_1) = q_1(t_1) = \pi. \end{cases}$$

We denote the energy of $\bar{q}$ by $\bar{e}$. Since $t_1 - t_{-1} \succeq |k_n|^{d-1}$, by the deduction as similar as the one in Lemma 2.17, we have

$$0 < \bar{e} \le \delta_n \exp\left(-\frac{C|k_n|^{\epsilon}}{\sqrt{\delta_n}}\right). \tag{2.2.37}$$

We set

$$e(t) = \frac{1}{2}|\dot{q}_1(t)|^2 - \delta_n(1 - \cos(q_1(t))),$$

it is easy to see that there exists $\bar{t} \in [t_{-1}, t_1]$ such that $e(\bar{t}) = \bar{e}$. Indeed, without loss of generality, we assume by contradiction that $q_1(t)$ lie above $\bar{q}(t)$ in the phase plane for all $t \in [t_{-1}, t_1]$, namely $\dot{q}_1(t) > \dot{\bar{q}}(t)$ for all $t \in [t_{-1}, t_1]$, which is contradicted by the boundary conditions $\bar{q}(t_{-1}) = q_1(t_{-1})$ and $\bar{q}(t_1) = q_1(t_1)$. Hence, there exists $\bar{t} \in [t_{-1}, t_1]$ such that $q_1(\bar{t}) = \bar{q}(\bar{t})$ and $\dot{q}_1(\bar{t}) = \dot{\bar{q}}(\bar{t})$, moreover we have $e(\bar{t}) = \bar{e}$.

By Euler-Lagrange equation, we can estimate $\dot{e}(t)$. More precisely,

$$\dot{e}(t) \preceq \mu_n \delta_n \dot{q}_1(t).$$

Hence, integrating from $\bar{t}$ to t, we have

$$\sup_{t \in [t_{-1}, t_1]} |e(t) - \bar{e}| \le \gamma \mu_n \delta_n, \tag{2.2.38}$$

where γ is a positive constant independent of n.

We proceed the proof of (i) by the following three steps.

a) $\dot{q}_1(t_1) > 0$.

we assume by contradiction that $\dot{q}_1(t_1) \le 0$. $q_1(t_1) = \pi$ together with (2.2.38) implies that for n large enough, $\dot{q}_1(t_1) < 0$. Let us denote by $(t_1 - \Delta t, t_1)$ the maximal interval on the left of t_1 on which $q_1(t) \ge \pi$. Since $\dot{q}_1(t_1) < 0$, then $\Delta t > 0$. From $q_1(t_{-1}) = -\pi < \pi = q_1(t_1)$, it follows that $t_1 - \Delta t > t_{-1}$. Let

$$\tilde{q}(t) = \begin{cases} q_1(t) & t \in [t_{-1}, t_1 - \Delta t), \\ 2\pi - q_1(t) & t \in [t_1 - \Delta t, t_1]. \end{cases}$$

It is easy to see that $(\tilde{q}, Q)$ is still an action minimizing orbit on $[t_{-1}, t_1]$. By the definition of Δt, we have that $\tilde{q}(t_1 - \Delta t) = \pi$ and

$$\dot{\tilde{q}}((t_1 - \Delta t)-) \cdot \dot{\tilde{q}}((t_1 - \Delta t)+) \le 0,$$

where $(t_1 - \Delta t)-$ denotes t tends to $t_1 - \Delta t$ from the left side and $(t_1 - \Delta t)+$ denotes t tends to $t_1 - \Delta t$ from the right side. By Euler-Lagrange equation, we have that $\dot{\tilde{q}}(t)$ is continuous for $t \in [t_{-1}, t_1]$. Hence, $\dot{\tilde{q}}(t_1 - \Delta t) = 0$. On the other hand, from (2.2.37) and (2.2.38), it follows that for n large enough, $\dot{\tilde{q}}(t_1 - \Delta t) \ne 0$. Therefore, we have $\dot{q}_1(t_1) > 0$.

b) $\dot{q}_1(t) > 0$ for $t \in \left[t_1 - \frac{\tau}{\delta'_n}, t_1\right]$.

Let $(\tilde{t}, t_1]$ be the maximal interval to the left of t_1 on which $\dot{q}_1 > 0$, hence we can denote the inverse function by $t(q_1)$. Let $\Gamma = \gamma\mu_n\delta_n$. It follows from (2.2.38) that for $q_1 < \pi$,

$$\begin{aligned} t_1 + \int_\pi^{q_1} \frac{ds}{\sqrt{2(\bar{e} - \Gamma + \delta_n(1-\cos(s)))}} &\le t(q_1), \\ &\le t_1 + \int_\pi^{q_1} \frac{ds}{\sqrt{2(\bar{e} + \Gamma + \delta_n(1-\cos(s)))}}. \end{aligned}$$

On the other hand, denoting the inverse function of $\hat{q}_1(t - t_1)$ by $\hat{t}(q_1)$, we have

$$\hat{t}(q_1) = t_1 + \int_\pi^{q_1} \frac{ds}{\sqrt{2(\delta_n(1-\cos(s)))}}.$$

Moreover, a simple calculation implies

$$|t(q_1) - \hat{t}(q_1)| \preceq \frac{\bar{e} + \Gamma}{(\delta_n(1-\cos(q_1)))^{\frac{3}{2}}},$$

which together with (2.2.26) implies that

$$|t(q_1) - \hat{t}(q_1)| \preceq \frac{\Gamma}{(\delta_n(1-\cos(q_1)))^{\frac{3}{2}}}. \tag{2.2.39}$$

It is easy to see that

$$|\dot{q}_1| + |\dot{\hat{q}}_1| \preceq \sqrt{\delta_n}. \tag{2.2.40}$$

Let

$$F(t) = \min\{1 - \cos(q_1(t)), 1 - \cos(\hat{q}_1(t - t_1))\}.$$

It follows from (2.2.39) and (2.2.40) that for $t \in [\tilde{t}, t_2]$

$$|q_1(t) - \hat{q}_1(t - t_1)| \preceq \frac{\Gamma\sqrt{\delta_n}}{(\delta_n F(t))^{\frac{3}{2}}}. \tag{2.2.41}$$

Hence, we have

$$\tilde{t} \le t_1 - \frac{\tau}{\delta'_n}.$$

In fact, we assume by contradiction that $\tilde{t} > t_1 - \frac{\tau}{\delta'_n}$. It follows from (2.2.41) that for n large enough,

$$q_1(\tilde{t}) \ge \frac{1}{2}\hat{q}_1\left(-\frac{\tau}{\delta'_n}\right). \tag{2.2.42}$$

If there exists $t^* \in [\tilde{t}, t_1]$ such that $\dot{q}_1(t^*) = 0$, then

$$|e(t^*)| \ge \delta_n \exp\left(-\frac{C|k_n|^{\frac{3\epsilon}{4}}}{\sqrt{\delta_n}}\right),$$

which is contradicted by (2.2.38) and (2.2.26). Since $\dot{q}_1(\tilde{t}) \geq 0$, we get $\dot{q}_1(\tilde{t}) > 0$, contradicted by the maximality of $[\tilde{t}, t_1]$. Therefore, we have $\tilde{t} \leq t_1 - \frac{\tau}{\delta'_n}$. i.e. $\dot{q}_1(t) > 0$ for $t \in \left[t_1 - \frac{\tau}{\delta'_n}, t_1\right]$.

c) $q_1(t_1 - \frac{\tau}{\delta'_n}) > 0$.

From (2.2.41) and (2.2.42), we have that for $t \in \left[t_1 - \frac{\tau}{\delta'_n}, t_1\right]$

$$|q_1(t) - \hat{q}_1(t - t_1)| \preceq \Gamma \delta_n^{-1} \exp\left(\frac{C|k_n|^{\frac{3\epsilon}{4}}}{\sqrt{\delta_n}}\right). \tag{2.2.43}$$

In terms of the definition of $\hat{q}_1(t)$, it follows from (2.2.26) that for n large enough,

$$\frac{1}{2}\hat{q}_1\left(-\frac{\tau}{\delta'_n}\right) \leq q_1\left(t_1 - \frac{\tau}{\delta'_n}\right) \leq \frac{3}{2}\hat{q}_1\left(-\frac{\tau}{\delta'_n}\right). \tag{2.2.44}$$

Hence, $q_1(t_1 - \frac{\tau}{\delta'_n}) > 0$, which together with $\dot{q}_1(t) > 0$ on $\left[t_1 - \frac{\tau}{\delta'_n}, t_1\right]$ implies $t_0 \leq t_1 - \frac{\tau}{\delta'_n}$.

Similarly, we have $t_1 + \frac{\tau}{\delta'_n} \leq t_2$. The proof of (i) is completed.

2.2.3.3.2. Proof of (ii)

We let $\Omega = [t_1 - \frac{\tau}{\delta'_n}, t_1 + \frac{\tau}{\delta'_n}]$. Since

$$\begin{aligned}
&\left|\delta_n \int_{t_0}^{t_2} (1 - \cos(q_1(t))) \cos(q_2(t)) dt - M_n(\bar{\omega}_2, Q(t_1), t_1)\right| \\
&\quad \leq C_1 \delta_n \left(\int_\Omega |\cos(q_2(t)) - \cos\left(\hat{q}_2(t) + q_2(t_1) - \hat{q}_2(t_1)\right)| dt \right. \\
&\quad\quad + \int_\Omega |\cos(q_1(t)) - \cos(\hat{q}_1(t - t_1))| dt + \int_{[t_0, t_2] \setminus \Omega} |1 - \cos(q_1(t))| dt \\
&\quad\quad \left. + \int_{\mathbb{R} \setminus \Omega} |1 - \cos(\hat{q}_1(t - t_1))| dt \right),
\end{aligned}$$

where $\hat{q}_2(t) = q_2(t_0) + \omega_2(t - t_0)$. Hence, it suffices to prove the following estimates

$$\delta_n \int_\Omega |\cos(q_2(t)) - \cos\left(\hat{q}_2(t) + q_2(t_1) - \hat{q}_2(t_1)\right)| dt \preceq \epsilon_n, \tag{2.2.45}$$

$$\delta_n \int_\Omega |\cos(q_1(t)) - \cos(\hat{q}_1(t - t_1))| dt \preceq \epsilon_n, \tag{2.2.46}$$

$$\delta_n \int_{[t_0, t_2] \setminus \Omega} |1 - \cos(q_1(t))| dt \preceq \epsilon_n, \tag{2.2.47}$$

$$\delta_n \int_{\mathbb{R} \setminus \Omega} |1 - \cos(\hat{q}_1(t - t_1))| dt \preceq \epsilon_n, \tag{2.2.48}$$

where we set $\epsilon_n = \sqrt{\delta_n} \exp(-\frac{\lambda}{\sqrt{\delta_n}})$. We will prove (2.2.45)-(2.2.48) in the following.

First of all, we prove (2.2.45). We set

$$\bar{q}(t)=\begin{cases}\hat{q}_1(t-t_1)\phi\left(t-(t_1-\frac{\tau}{\delta_n'})\right), & t_0\le t\le t_1,\\ \left(\hat{q}_1(t-t_1)-2\pi\right)\phi\left(-t+(t_1+\frac{\tau}{\delta_n'})\right)+2\pi, & t_1\le t\le t_2,\end{cases}$$

where ϕ is a C^∞ function as follow

$$\phi(t)=\begin{cases}0, & t\le 0,\\ 1, & t\ge 1.\end{cases}$$

We set $\hat{Q}(t)=Q(t_0)+\bar{\omega}(t-t_0)$, hence $(\bar{q},\hat{Q})(t_0)=(q_1,Q)(t_0)$ and $(\bar{q},\hat{Q})(t_2)=(q_1,Q)(t_2)$. From the minimality of (q_1,Q), it follows that

$$\int_{t_0}^{t_2}L_n\left(q_1,Q,\dot{q}_1,\dot{Q}\right)dt\le\int_{t_0}^{t_2}L_n\left(\bar{q},\hat{Q},\dot{\bar{q}},\dot{\hat{Q}}\right)dt.$$

Since $t_2-t_0\ge 2\tau/\delta_n'$, we let $\bar{\Omega}=[t_1-\frac{\tau}{\delta_n'}+1,t_1+\frac{\tau}{\delta_n'}-1]$, then (2.2.31) and a direct calculation implies

$$\begin{aligned}\int_{t_0}^{t_2}(1-\cos(\bar{q}(t)))dt&=\int_{\bar{\Omega}}1-\cos(\hat{q}_1(t-t_1))dt\\&\quad+\int_{[t_0,t_2]\setminus\bar{\Omega}}(1-\cos(\bar{q}(t)))dt,\\&\preceq\frac{1}{\sqrt{\delta_n}}\int_0^{2\pi}\sqrt{1-\cos(\hat{q}_1)}d\hat{q}_1,\\&\preceq\frac{1}{\sqrt{\delta_n}},\end{aligned}$$

hence,

$$\int_{t_0}^{t_2}(1-\cos(\bar{q}))dt\preceq\frac{1}{\sqrt{\delta_n}}. \tag{2.2.49}$$

Moreover, let $\bar{\omega}=(\bar{\omega}_2,\dots,\bar{\omega}_d)$, in terms of the definition of L_n as (2.2.27), we have

$$\begin{aligned}&\int_{t_0}^{t_2}L_n\left(\bar{q},\hat{Q},\dot{\bar{q}},\dot{\hat{Q}}\right)dt\\&\quad\le\int_{t_0}^{t_2}\frac{1}{2}\sum_{i=3}^{d}\frac{|\bar{\omega}_i|^2}{|l_{ni}|^2}+\frac{|\bar{\omega}_2|^2}{2|k_n'|^2}\\&\qquad+\frac{1}{|k_n|^2}\left(\frac{1}{2}|\dot{\bar{q}}|^2+\delta_n(1-\cos(\bar{q}))\right)dt+\frac{1}{|k_n|^2}\mu_n\delta_n\int_{t_0}^{t_2}(1-\cos(\bar{q}))dt,\end{aligned}$$

hence, from (2.2.49), it follows that

$$\begin{aligned}&\int_{t_0}^{t_2}L_n\left(\bar{q},\hat{Q},\dot{\bar{q}},\dot{\hat{Q}}\right)dt\\&\quad\le\int_{t_0}^{t_2}\frac{1}{2}\sum_{i=3}^{d}\frac{|\bar{\omega}_i|^2}{|l_{ni}|^2}+\frac{|\bar{\omega}_2|^2}{2|k_n'|^2}\\&\qquad+\frac{1}{|k_n|^2}\left(\frac{1}{2}|\dot{\bar{q}}|^2+\delta_n(1-\cos(\bar{q}))\right)dt+\frac{C}{|k_n|^2}\mu_n\sqrt{\delta_n}.\end{aligned}$$

Since μ_n is small enough for n large enough, we have

$$\begin{aligned}&\int_{t_0}^{t_2} L_n\left(q_1, Q, \dot{q}_1, \dot{Q}\right) dt\\ &\qquad \geq \int_{t_0}^{t_2} \frac{1}{2}\sum_{i=3}^{d}\frac{|\dot{q}_i|^2}{|l_{ni}|^2} + \frac{|\dot{q}_2|^2}{2|k_n'|^2} dt + \frac{1}{2}\frac{1}{|k_n|^2}\int_{t_0}^{t_2}(|\dot{q}_1|^2 + \delta_n(1-\cos(q_1)))dt.\end{aligned}$$

It is easy to see that

$$\frac{1}{2}|\dot{q}_i|^2 = \frac{1}{2}|\bar{\omega}_i|^2 + \langle \bar{\omega}_i, \dot{q}_i - \bar{\omega}_i\rangle + \frac{1}{2}|\dot{q}_i - \bar{\omega}_i|^2, \quad \text{for} \quad i = 2, \dots, d,$$

and the mean value of $\dot{q}_i - \bar{\omega}_i$ vanishes on $[t_0, t_2]$ for $i = 2, \dots, d$. Hence,

$$\begin{aligned}&\int_{t_0}^{t_2} L_n\left(q_1, Q, \dot{q}_1, \dot{Q}\right) dt\\ &\qquad \geq \int_{t_1}^{t_3} \frac{1}{2}\sum_{i=3}^{d}\frac{|\bar{\omega}_i|^2}{|l_{ni}|^2} + \frac{|\bar{\omega}_2|^2}{2|k_n'|^2} dt + \frac{1}{2}\frac{1}{|k_n|^2}\int_{t_0}^{t_2}(|\dot{q}_1|^2 + \delta_n(1-\cos(q_1)))dt.\end{aligned}$$

From the deduction above, we have

$$\frac{1}{2}\int_{t_0}^{t_2}(|\dot{q}_1|^2 + \delta_n(1-\cos(q_1)))dt \leq \int_{t_0}^{t_2}\left(\frac{1}{2}|\dot{\bar{q}}|^2 + \delta_n(1-\cos(\bar{q}))\right)dt + C\mu_n\sqrt{\delta_n}.$$

From the definition of $\bar{q}$ and (2.2.31), a direct calculation implies

$$\begin{aligned}&\int_{t_0}^{t_2}\left(\frac{1}{2}|\dot{\bar{q}}|^2 + \delta_n(1-\cos(\bar{q}))\right)dt\\ &\quad = \int_{\bar{\Omega}}\left(\frac{1}{2}|\dot{\hat{q}}_1(t-t_2)|^2 + \delta_n(1-\cos(\hat{q}_1(t-t_2))\right)dt\\ &\qquad + \int_{[t_0,t_2]\setminus\bar{\Omega}}(1-\cos(\bar{q}(t)))dt,\\ &\quad \preceq \sqrt{\delta_n}\int_0^{2\pi}\sqrt{1-\cos(\hat{q}_1)}d\hat{q}_1,\\ &\quad \preceq \sqrt{\delta_n},\end{aligned}$$

hence,

$$\frac{1}{2}\int_{t_0}^{t_2}(|\dot{q}_1|^2 + \delta_n(1-\cos(q_1)))dt \preceq \sqrt{\delta_n}. \tag{2.2.50}$$

From Euler-Lagrange equation on (q_1, Q), it follows that

$$|\ddot{Q}| \leq \mu_n\delta_n(1-\cos(q_1)).$$

Hence,

$$|\dot{Q}(t) - \dot{Q}(t')| \leq \mu_n\delta_n\int_{t'}^{t}(1-\cos(q_1))dt, \quad \text{for any } t,\ t' \in [t_0, t_2].$$

By (2.2.50), we have

$$|\dot{Q}(t)-\dot{Q}(t')|\preceq \mu_n\sqrt{\delta_n},\quad \text{for any } t,\ t'\in[t_0,t_2].$$

Moreover,

$$|\dot{Q}(t)-\bar{\omega}|\preceq \mu_n\sqrt{\delta_n},\quad \text{for any } t\in[t_0,t_2],$$

which yields

$$\sup_{\Omega}|Q(t)-(\hat{Q}(t)+Q(t_1)-\hat{Q}(t_1))|\preceq \frac{|k_n|^2\mu_n}{\sqrt{\delta_n}},$$

if we integrate $\dot{Q}(t)$ from t_1 to t. In particular, we have

$$\sup_{\Omega}|q_2(t)-(\hat{q}_2(t)+q_2(t_1)-\hat{q}_2(t_1))|\preceq \frac{|k_n|^2\mu_n}{\sqrt{\delta_n}}.$$

Since

$$|\cos(\theta_1)-\cos(\theta_2)|\le|\theta_1-\theta_2|, \tag{2.2.51}$$

then the proof of (2.2.45) is completed if we take n large enough.

Second, we prove (2.2.46). By (2.2.44), we have

$$\frac{1}{2}\hat{q}_1\left(-\frac{\tau}{\delta_n'}\right)\le q_1\left(t_1-\frac{\tau}{\delta_n'}\right)\le\frac{3}{2}\hat{q}_1\left(-\frac{\tau}{\delta_n'}\right),$$

hence, from (2.2.31), it follows that

$$q_1\left(t_1-\frac{\tau}{\delta_n'}\right)\succeq\exp\left(-\frac{|k_n|^{\frac{3\epsilon}{4}}\tau}{\sqrt{\delta_n}}\right).$$

Moreover, the above inequality and (2.2.43) imply that for n large enough and $t\in(t_1-\frac{\tau}{\delta_n'},t_2]$,

$$|q_1(t)-\hat{q}_1(t-t_1)|\preceq\frac{\Gamma}{\delta_n(1-\cos(\hat{q}_1(t-t_1)))^{\frac{3}{2}}}\preceq\epsilon_n.$$

The case in which $t\in[t_1,t_1+\frac{\tau}{\delta_n'}]$ is similar, more precisely,

$$|q_1(t)-\hat{q}_1(t-t_1)|\preceq\epsilon_n,\quad\text{for}\quad t\in\left[t_1,t_1+\frac{\tau}{\delta_n'}\right].$$

Therefore, from (2.2.51), it follows that

$$|\cos(q_1(t))-\cos(\hat{q}_1(t-t_1))|\preceq\epsilon_n,\quad\text{for}\quad t\in\Omega,$$

hence, we complete the proof of (2.2.46).

Third, we prove (2.2.47). It suffices to prove for $\Omega'=\left[t_{-1}+\frac{\tau}{\delta_n'},t_2-\frac{\tau}{\delta_n'}\right]$

$$\delta_n\int_{\Omega'}|1-\cos(q_1(t))|dt\preceq\epsilon_n,$$

where t_{-1} satisfies $q_1(t_{-1})=-\pi$.

Let $\tilde{q}_1$ be the solution of A_n satisfying boundary conditions

$$\begin{cases} \tilde{q}_1\left(t_{-1}+\frac{\tau}{\delta_n'}\right)=q_1\left(t_{-1}+\frac{\tau}{\delta_n'}\right), \\ \\ \tilde{q}_1\left(t_1-\frac{\tau}{\delta_n'}\right)=q_1\left(t_1-\frac{\tau}{\delta_n'}\right). \end{cases}$$

From the minimality of (q_1, Q), it follows that

$$\int_{\Omega'} L_n\left(q_1, Q, \dot{q}_1, \dot{Q}\right) dt \le \int_{\Omega'} L_n\left(\tilde{q}_1, Q, \dot{\tilde{q}}_1, \dot{Q}\right) dt.$$

By (2.2.27), we have

$$\begin{aligned} &\int_{\Omega'} \frac{1}{2}|\dot{q}_1|^2+\delta_n(1-\cos(q_1))+\mu_n\delta_n(1-\cos(q_1))\cos(q_2)dt \\ &\qquad \le \int_{\Omega'} \frac{1}{2}|\dot{\tilde{q}}_1|^2+\delta_n(1-\cos(\tilde{q}_1))+\mu_n\delta_n(1-\cos(\tilde{q}_1))\cos(q_2)dt. \end{aligned}$$

From the construction of $\tilde{q}_1$, let $\tilde{e}$ be the energy of $\tilde{q}_1$, a similar argument as the one in Lemma 2.17 implies

$$0<\tilde{e}\le \delta_n \exp\left(-\frac{C|k_n|^\epsilon}{\sqrt{\delta_n}}\right).$$

Based on the change of variable $dq=\dot{q}dt$, a direct calculation gives

$$\begin{aligned} &\int_{\Omega'} \frac{1}{2}|\dot{\tilde{q}}_1|^2+\delta_n(1-\cos(\tilde{q}_1))+\mu_n\delta_n(1-\cos(\tilde{q}_1))\cos(q_2)dt \\ &\qquad \preceq \exp\left(-\frac{C|k_n|^\epsilon}{\sqrt{\delta_n}}\right). \end{aligned}$$

For n large enough, we have

$$\begin{aligned} &\frac{1}{2}\int_{\Omega'} |\dot{q}_1|^2+\delta_n(1-\cos(q_1))dt \\ &\qquad \le \int_{\Omega'} \frac{1}{2}|\dot{q}_1|^2+\delta_n(1-\cos(q_1))+\mu_n\delta_n(1-\cos(q_1))\cos(q_2)dt, \end{aligned}$$

which together with $|\dot{q}_1|\ge 0$ implies

$$\delta_n\int_{\Omega'} |1-\cos(q_1(t))|dt \le \int_{\Omega'} |\dot{q}_1|^2+\delta_n(1-\cos(q_1))dt \preceq \epsilon_n,$$

hence, (2.2.47) is proved.

Finally, the inequality (2.2.48) can be obtained by a direct calculation. More precisely,

$$\begin{aligned} \delta_n\int_{\mathbb{R}\setminus\Omega} |1-\cos(\hat{q}_1(t-t_1))|dt =&\delta_n\int_{-\infty}^{t_1-\frac{\tau}{\delta_n'}} 1-\cos(\hat{q}_1(t-t_1))dt \\ &+\delta_n\int_{t_1+\frac{\tau}{\delta_n'}}^{\infty} 1-\cos(\hat{q}_1(t-t_1))dt. \end{aligned}$$

By (2.2.31), $\hat{q}_1(t) = 4\arctan\big(\exp(\sqrt{\delta_n}(t-t_1))\big)$, we have

$$\begin{aligned}\delta_n\int_{-\infty}^{t_1-\frac{\tau}{\delta_n'}} 1-&\cos(\hat{q}_1(t-t_1))dt\\ &\preceq \delta_n\int_{-\infty}^{t_1-\frac{\tau}{\delta_n'}}\arctan^2\Big(\exp(\sqrt{\delta_n}(t-t_1))\Big)\,dt,\\ &\preceq \delta_n\int_{-\infty}^{t_1-\frac{\tau}{\delta_n'}}\exp\Big(2\sqrt{\delta_n}(t-t_1)\Big)\,dt,\\ &\preceq \sqrt{\delta_n}\exp\left(-\frac{2|k_n|^{\frac{3\epsilon}{4}}\tau}{\sqrt{\delta_n}}\right).\end{aligned}$$

It is similar to get

$$\delta_n\int_{t_1+\frac{\tau}{\delta_n'}}^{\infty} 1-\cos(\hat{q}_1(t-t_1))dt \preceq \sqrt{\delta_n}\exp\left(-\frac{2|k_n|^{\frac{3\epsilon}{4}}\tau}{\sqrt{\delta_n}}\right).$$

Hence, (2.2.48) is proved for n large enough,.

So far, we complete the proof of (2.2.45)-(2.2.48) and also the proof of Lemma 2.24. ☐

Remark 2.25 *In the proof of (2.2.45), as a bonus we obtain an important estimate on* $\dot{Q}(t)$. *Namely, let* $(q_1(t), Q(t))$ *be the action minimizing orbit of* L_n, *then for any* $t', t'' \in \mathbb{R}$ *and* $t \in [t', t'']$ *we have*

$$\left|\dot{Q}(t) - \frac{Q(t'')-Q(t')}{t''-t'}\right| \le C\mu_n\sqrt{\delta_n}.$$

From Remark 2.25 and the definition of Melnikov function (2.2.32), it follows that for n large enough,

$$|M_n(\omega_2, Q_1, t_1) - M_n(\bar{\omega}_2, Q_1, t_1)| \le C\sqrt{\delta_n}\exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right), \tag{2.2.52}$$

where λ is the same as the one in (2.2.34).

2.2.3.4. Proof of Theorem 2.22

Based on the minimal property of the orbits on an invariant Lagrangian torus and its graph property, passing through each $x \in \mathbb{T}^d$, there is a unique minimal curve $q(t)$ with rotation vector ω if the Hamiltonian flow generated by H_n admits a Lagrangian torus with rotation vector ω. Hence, it is sufficient to prove the existence of some point in $\mathbb{T}^d$ where no minimal curve passes through.

Indeed, any minimal curve does not pass through the subspace $(\pi, 0) \times \mathbb{T}^{d-2}$. It implies Lemma 2.24. Let us assume contrary, namely, there exists $\bar{t}_1$ such that

$$q_1(\bar{t}_1) = \pi, \quad q_2(\bar{t}_1) = 0,$$

where $q(t) = (q_1, q_2, \ldots, q_d)(t)$ is a minimal orbit in the universal covering space $\mathbb{R}^d$. Because of $\omega_1 \neq 0$, there exist t_0 and t_1 such that

$$q_1(t_0) = 0, \quad q_1(t_2) = 2\pi.$$

Obviously, $t_0 < \bar{t}_1 < t_2$. From (2.2.1), we have

$$t_2 - t_0 \geq \frac{2\pi}{C_0} |k_n|^{d-1},$$

where C_0 is the constant in (2.2.1). Let $\tilde{t}_1$ be the last time before $\bar{t}_1$ or the first time after $\bar{t}_1$ such that

$$|q_2(\tilde{t}_1) - q_2(\bar{t}_1)| = \pi.$$

By (2.2.22), we have

$$|\tilde{t}_1 - \bar{t}_1| \sim \frac{1}{|\omega_2|} \leq C_1.$$

Without loss of generality, one can assume $\omega_1 > 0$ and $\omega_2 > 0$. Consider a solution $\tilde{q}_1$ of A_n on $(t_0, \tilde{t}_1)$ and on $(\tilde{t}_1, t_2)$ with boundary conditions respectively

$$\begin{cases} \tilde{q}_1(t_0) = q_1(t_0) = 0, \\ \tilde{q}_1(\tilde{t}_1) = q_1(\bar{t}_1) = \pi, \end{cases} \qquad \begin{cases} \tilde{q}_1(\tilde{t}_1) = q_1(\bar{t}_1) = \pi, \\ \tilde{q}_1(t_2) = q_1(t_2) = 2\pi. \end{cases}$$

Since $q = (q_1, q_2, \dots, q_d)$ is assumed to be a minimal curve, setting $Q = (q_2, \dots, q_d)$, we have

$$\int_{t_0}^{t_2} L_n(\tilde{q}_1, Q, \dot{\tilde{q}}_1, \dot{Q}) dt - \int_{t_0}^{t_2} L_n(q_1, Q, \dot{q}_1, \dot{Q}) dt \geq 0. \tag{2.2.53}$$

See Fig.3, where $x_1 = (q_1(\bar{t}_1), q_2(\bar{t}_1)) = (\pi, 0)$, $x_0 = (q_1(t_0), q_2(t_0)) = (0, q_2(t_0))$, $x_2 = (q_1(t_2), q_2(t_2)) = (2\pi, q_2(t_2))$, $\tilde{x}_1' = (\pi, -\pi)$ and $\tilde{x}_1'' = (\pi, \pi)$.



Figure 3: The projections of the curves $(q_1(t), Q(t))$ and $(\tilde{q}_1(t), Q(t))$ on $[0, 2\pi] \times \mathbb{R}$

$(\tilde{q}_1(t), q_2(t))$ passes through the point $\tilde{x}_1'$ or $\tilde{x}_1''$. Thus, by the construction of L_n, we obtain from (2.2.53) that

$$\begin{aligned} &\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1) dt - \int_{t_0}^{t_2} A_n(q_1, \dot{q}_1) dt \\ &\qquad \geq \mu_n \left(\int_{t_0}^{t_2} \tilde{P}_n(q_1, Q) dt - \int_{t_0}^{t_2} \tilde{P}_n(\tilde{q}_1, Q) dt \right), \end{aligned} \tag{2.2.54}$$

where

$$\tilde{P}_n(q_1, Q) = \mu_n \delta_n (1 - \cos q_1) \cos q_2.$$

By Lemma 2.24, we have

$$\begin{aligned} &\int_{t_0}^{t_2} \tilde{P}_n(\tilde{q}_1, Q) dt - \int_{t_0}^{t_2} \tilde{P}_n(q_1, Q) dt \\ &\qquad \leq M_n(\bar{\omega}_2, Q(\tilde{t}_1), \tilde{t}_1) - M_n(\bar{\omega}_2, Q(\bar{t}_1), \bar{t}_1) + C\sqrt{\delta_n} \exp\left(-\frac{\lambda}{\sqrt{\delta_n}} \right), \end{aligned}$$

where $\bar{\omega} = (Q(t_2) - Q(t_0))/(t_2 - t_0)$. Here, the approximation from $M_n(\bar{\omega}_2, Q(\tilde{t}_1), \tilde{t}_1)$ to $\int_{t_0}^{t_2} \tilde{P}_n(\tilde{q}_1, Q)dt$ can not be obtained directly by Lemma 2.24, since $(\tilde{q}_1, Q(t))$ may be not minimal. But based on the simplicity of $\tilde{q}_1$, a much simpler calculation than the one in Lemma 2.24 implies that the approximation in Lemma 2.24 is still verified for the orbit $(\tilde{q}_1, Q(t))$. We omit it.

By Lemma 2.23 and (2.2.52), we have

$$\int_{t_0}^{t_2} \tilde{P}_n(\tilde{q}_1, Q)dt - \int_{t_0}^{t_2} \tilde{P}_n(q_1, Q)dt \preceq -\exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right).$$

From (2.2.54), it follows that

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt \succeq \mu_n \exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right). \tag{2.2.55}$$

On the other hand, consider a solution $\bar{q}_1$ of A_n on $(t_0, \bar{t}_1)$ and on $(\bar{t}_1, t_2)$ with boundary conditions respectively

$$\begin{cases} \bar{q}_1(t_0) = q_1(t_0) = 0, \\ \bar{q}_1(\bar{t}_1) = q_1(\bar{t}_1) = \pi, \end{cases} \qquad \begin{cases} \bar{q}_1(\bar{t}_1) = q_1(\bar{t}_1) = \pi, \\ \bar{q}_1(t_2) = q_1(t_2) = 2\pi. \end{cases}$$

We have

$$\int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt \geq \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt.$$

Since $|\tilde{t}_1 - \bar{t}_1| \leq C_1$, An argument similar as the one to Fig. 2 implies

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt \preceq |k_n|^{d-1}\delta_n \exp\left(-\frac{C|k_n|^{\epsilon}}{\sqrt{\delta_n}}\right).$$

From (2.2.26),

$$\mu_n \sim \exp\left(-|k_n|^{\frac{d}{2}-\frac{1}{2}+\frac{\epsilon}{3}}\right),$$

it follows that

$$\int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt \preceq \frac{\mu_n}{|k_n|} \exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right).$$

Hence, for any $\bar{t}_1 \in (t_0, t_2)$, we can find $\tilde{t}_1$ such that

$$\begin{aligned} \int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(q_1, \dot{q}_1)dt &\leq \int_{t_0}^{t_2} A_n(\tilde{q}_1, \dot{\tilde{q}}_1)dt - \int_{t_0}^{t_2} A_n(\bar{q}_1, \dot{\bar{q}}_1)dt, \\ &\preceq \frac{\mu_n}{|k_n|} \exp\left(-\frac{\lambda}{\sqrt{\delta_n}}\right), \end{aligned}$$

which is contradicted by (2.2.55) for large n. this completes the proof of Theorem 2.22. □

3. DESTRUCTION OF ALL LAGRANGIAN TORI

In this section, we are concerned with Lagrangian tori as follow:

Definition 3.1 *$\mathcal{T}^d$ is called d dimensional Lagrangian torus with the rotation vector ω if*

- *$\mathcal{T}^d$ is a Lagrangian submanifold;*
- *$\mathcal{T}^d$ is invariant for the Hamiltonian flow Φ_H^t generated by H.*

For positive definite Hamiltonian systems, if a Lagrangian torus is invariant under the Hamiltonian flow, it is then the graph over $\mathbb{T}^d$ (see [BP]).

In the following subsections, we consider the destruction of all Lagrangian tori of symplectic twist maps. Based on the correspondence between symplectic twist maps and Hamiltonian systems, it can be achieved to destruct all Lagrangian tori of Hamiltonian systems.

3.1. A toy model

To show the basic ideas, we are beginning with a toy model whose generating function is as follow:

$$h_n(x,x') = h_0(x,x') - \frac{5}{4n^2}\sin(nx') - \frac{1}{16n^2}\cos(2nx'), \tag{3.1.1}$$

where $h_0(x,x') = \frac{1}{2}(x-x')^2$. Let $f_n(x,y) = (x',y')$ be the exact area-preserving twist map generated by (3.1.1), then

$$\begin{cases} y = -\partial_1 h_n(x,x') = x' - x, \\ y' = \partial_2 h_n(x,x') = x' - x - \frac{5}{4n}\cos(nx') + \frac{1}{8n}\sin(2nx'). \end{cases}$$

We set $\phi_n(x) = -\frac{5}{4n}\cos(nx) + \frac{1}{8n}\sin(2nx)$, then

$$f_n(x,y) = (x+y, y + \phi_n(x+y)). \tag{3.1.2}$$

In [H2], Herman found a criterion of total destruction of invariant circles. By Birkhoff graph theorem (see [H4]), if f_n admits an invariant circle, then the invariant circle is a Lipschitz graph. We denote the graph by ψ_n, then it follows from [H5] that

$$\psi_n \circ g_n - \psi_n = \phi_n \circ g_n,$$

where $g_n = \mathrm{Id} + \psi_n$. This is equivalent to

$$\frac{1}{2}(g_n + g_n^{-1}) = \mathrm{Id} + \frac{1}{2}\phi_n. \tag{3.1.3}$$

Let $\mathfrak{D}_n$ be the set of differentiable points of g_n, then $\mathfrak{D}_n$ has full Lebesgue measure on $\mathbb{R}$ since g_n is a Lipschitz function. For $x \in \mathfrak{D}_n$, we differentiae (3.1.3),

$$\frac{1}{2}(Dg_n(x) + (Dg_n)^{-1}(g_n^{-1}(x))) = 1 + \frac{1}{2}D\phi_n(x).$$

Let $G_n = ||Dg_n||_{L^\infty}$. It is easy to see that for $\varepsilon > 0$, there exists $\tilde{x} \in \mathfrak{D}_n$ such that $Dg_n(\tilde{x}) \geq G_n - \varepsilon$. Let $M_n = \max D\phi_n$, we have

$$\frac{1}{2}\left(G_n + \frac{1}{G_n} - \varepsilon\right) \leq 1 + \frac{1}{2}M_n.$$

Since $\varepsilon > 0$ is arbitrarily small, then

$$\frac{1}{2}\left(G_n + \frac{1}{G_n}\right) \leq 1 + \frac{1}{2}M_n.$$

Hence,

$$G_n \leq 1 + \frac{1}{2}M_n + \left(M_n + \frac{1}{4}{M_n}^2\right)^{\frac{1}{2}}. \tag{3.1.4}$$

Obviously, for $x \in \mathfrak{D}_n$, we have

$$\frac{1}{G_n} \leq 1 + \frac{1}{2}D\phi_n(x).$$

Let $m_n = \min D\phi_n$, then we have

$$\frac{1}{G_n} \leq 1 + \frac{1}{2}m_n,$$

which together with (3.1.4) implies that

$$\frac{1}{1+\frac{1}{2}m_n} \leq 1 + \frac{1}{2}M_n + \left(M_n + \frac{1}{4}{M_n}^2\right)^{\frac{1}{2}}.$$

Therefore, it is sufficient for total destruction of invariant circles to construct $\phi_n(x)$ such that

$$\frac{1}{1+\frac{1}{2}\min D\phi_n} > 1 + \frac{1}{2}\max D\phi_n + \left(\max D\phi_n + \frac{1}{4}(\max D\phi_n)^2\right)^{\frac{1}{2}}. \tag{3.1.5}$$

In our construction,

$$D\phi_n(x) = \frac{5}{4}\sin(nx) + \frac{1}{4}\cos(2nx). \tag{3.1.6}$$

A simple calculation implies

$$\begin{cases} \min D\phi_n(x) = -\frac{3}{2}, & \text{attained at } x = \frac{3}{2n} + \frac{2\pi k}{n}, \\ \max D\phi_n(x) = 1, & \text{attained at } x = \frac{\pi}{2n} + \frac{2\pi k}{n}, \end{cases}$$

where $k \in \mathbb{Z}$. Hence, (3.1.5) holds. Moreover, the exact area-preserving twist map generated by (3.1.1) admits no invariant circles.

By interpolation inequality ([H2]), for a small positive constant δ, we have

$$||\phi_n||_{C^{1-\delta}} \leq 2||\phi_n||_{C^0}^{\delta}||D\phi_n||_{C^0}^{1-\delta}.$$

From the construction of ϕ_n, it follows that $||\phi_n||_{C^0} \to 0$, as $n \to \infty$ and $||D\phi_n||_{C^0}$ is bounded. Hence,

$$||\phi_n||_{C^{1-\delta}} \to 0 \quad \text{as} \quad n \to \infty,$$

which implies that

$$||h_n - h_0||_{C^{2-\delta}} \to 0 \quad \text{as} \quad n \to \infty.$$

3.2. C^∞ case

In [H5], Herman extended the criterion (3.1.5) to multi-degrees of freedom. More precisely, for exact symplectic twist map of $\mathrm{T}^*\mathbb{T}^d$ of the following form

$$f(x,y) = (x+y, y + d\Psi(x+y)), \tag{3.2.1}$$

where $\Psi \in C^r(\mathbb{T}^d, \mathbb{R})$, $r \geq 2$ and

$$d\Psi = \left(\frac{\partial \Psi}{\partial x_1}, \cdots, \frac{\partial \Psi}{\partial x_d}\right).$$

Let $E(x)$ be the derivative matrix of $d\Psi$ and $T(x) = \frac{1}{d}\mathrm{tr}E(x)$, where tr denotes the trace of $E(x)$. From a similar argument as the deduction of (3.1.5), it follows that it is sufficient for total destruction of Lagrangian tori to construct $T(x)$ such that

$$\frac{1}{1+\frac{1}{2}\min T(x)} > 1 + \frac{1}{2}\max T(x) + \left(\max T(x) + \frac{1}{4}(\max T(x))^2\right)^{\frac{1}{2}}. \tag{3.2.2}$$

Moreover, for $T(x) \to 0$, (3.2.2) implies

$$-\frac{1}{2}\min T(x) > \sqrt{\max T(x)} + O(\max T(x)). \tag{3.2.3}$$

Herman constructed a sequence $\{\Psi_n\}_{n\in\mathbb{N}}$ that satisfies (3.2.3). It is easy to see $T_n(x) = \frac{1}{d}\Delta\Psi_n$ where Δ denotes the Laplacian. Since $T_n(x)$ is 2π-periodic, it is enough to construct it on $[-\pi,\pi]^d$. More precisely,

$$T_n(x) = \begin{cases} T_n^+(x), & x \in [0,\pi]^d, \\ T_n^-(x), & x \in [-\pi, 0]^d, \\ 0, & \text{others}. \end{cases}$$

where $T_n(x)$ is C^∞ function, $T_n^+(x)$ and $T_n^-(x)$ have the following forms respectively.

$T_n^+(x)$ satisfies:

$$\begin{cases} \operatorname{supp} T_n^+(x) \subset [0,\pi]^d, \\ \max T_n^+(x) = \frac{1}{n}, \end{cases}$$

$T_n^-(x)$ satisfies:

$$\begin{cases} \operatorname{supp} T_n^-(x) = B_{R_n}(x_0), \\ \max T_n^-(x) = \frac{1}{\sqrt{n}}, \\ R_n \sim \left(\frac{1}{\sqrt{n}}\right)^{\frac{1}{d}}, \\ x_0 = \left(-\frac{\pi}{2}, \cdots, -\frac{\pi}{2}\right), \end{cases}$$

where $f \sim g$ means that $\frac{1}{C}g < f < Cg$ for a constant $C > 1$. Hence, we obtain a sequence of $\{T_n(x)\}_{n\in\mathbb{N}}$ with bounded C^d norms and satisfying

$$\int_{\mathbb{T}^d} T_n(x)dx = 0.$$

Form interpolation inequality, it follows that $T_n(x) \to 0$ as $n \to \infty$ in the $C^{d-\delta}$ topology for any $\delta > 0$.

Let Ψ_n be the unique function in $C^\infty(\mathbb{T}^d,\mathbb{R})$ such that

$$\int_{\mathbb{T}^d}\Psi_n(x)dx = 0 \quad \text{and} \quad \frac{1}{d}\Delta\Psi_n(x) = T_n(x).$$

By Schauder estimates one knows that for any $\delta > 0$, $\Psi_n(x) \to 0$ as $n \to \infty$ in the $C^{d+2-\delta}$ topology. From the construction of $T_n(x)$, it is easy to see that (3.2.3) is verified.

Above all, we have the following theorem

Theorem 3.2 *All Lagrangian tori of an integrable symplectic twist map with $d \geq 1$ degrees of freedom can be destructed by C^∞ perturbations of the generating function and the perturbations are arbitrarily small in the $C^{d+2-\delta}$ topology for a small given constant $\delta > 0$.*

Based on the correspondence between symplectic twist maps and Hamiltonian systems, we have the following corollary.

Corollary 3.3 *All Lagrangian tori of an integrable Hamiltonian system with $d \geq 2$ degrees of freedom can be destructed by C^∞ perturbations which are arbitrarily small in the $C^{d+1-\delta}$ topology for a small given constant $\delta > 0$.*

3.3. An approximation lemma

In this subsection, we will prove a lemma on C^∞ functions approximated by trigonometric polynomials. First of all, we need some notations. Define

$$C^\infty_{2\pi}(\mathbb{R}^d,\mathbb{R}) := \left\{f : \mathbb{R}^d \to \mathbb{R} | f \in C^\infty(\mathbb{R}^d,\mathbb{R}) \text{ and } 2\pi - \text{periodic in } x_1,\dots,x_d\right\}.$$

Let $f(x) \in C^\infty_{2\pi}(\mathbb{R}^d,\mathbb{R})$. The m-th Fejér-polynomial of f with respect to x_j is given by

$$F^{[j]}_m(f)(x) := \frac{1}{m\pi}\int_{-\pi/2}^{\pi/2} f(x+2e_j)\left(\frac{\sin(mt)}{\sin t}\right)^2 dt, \tag{3.3.1}$$

where $x \in \mathbb{R}^d$, $m \in \mathbb{N}$, $j \in \{1,\dots,d\}$ and e_j is the j-th vector of the canonical basis of $\mathbb{R}^d$. $F^{[j]}_m(f)(x)$ is a trigonometric polynomial in x_j of degree at most $m-1$. By [Z],

$$\frac{1}{m\pi}\int_{-\pi/2}^{\pi/2}\left(\frac{\sin(mt)}{\sin t}\right)^2 dt = 1,$$

hence, from (3.3.1), we have

$$||F^{[j]}_m(f)||_{C^0} \leq ||f||_{C^0}.$$

We denote

$$P^{[j]}_m(f) := 2F^{[j]}_{2m}(f) - F^{[j]}_m(f).$$

It is easy to see that $P^{[j]}_m(f)$ is a trigonometric polynomial in x_j of degree at most $2m-1$. Moreover,

$$||P^{[j]}_m(f)||_{C^0} \leq 3||f||_{C^0}, \tag{3.3.2}$$

$$P_m^{[j]}(af+bg) = aP_m^{[j]}(f) + bP_m^{[j]}(g), \tag{3.3.3}$$

where $a, b \in \mathbb{R}$ and $f, g \in C^\infty_{2\pi}(\mathbb{R}^d, \mathbb{R})$. For $k \in \{1, \dots, d\}$, $j_1, \dots, j_k \in \{1, \dots, d\}$ with $j_p \neq j_q$ for $p \neq q$. Let $m_1, \dots, m_k \in \mathbb{N}$ and $f \in C^\infty_{2\pi}(\mathbb{R}^d, \mathbb{R})$, we define

$$P^{[j_1,\dots,j_k]}_{m_1,\dots,m_k}(f) := P^{[j_1]}_{m_1}\left(P^{[j_2]}_{m_2}\left(\cdots\left(P^{[j_k]}_{m_k}(f)\right)\cdots\right)\right). \tag{3.3.4}$$

It is easy to see that for all $l \in \{1, \dots, k\}$, $P^{[j_1,\dots,j_k]}_{m_1,\dots,m_k}(f)$ are trigonometric polynomials in x_l of degree at most $2m_l - 1$. We have the following lemma.

Lemma 3.4 *Let $f \in C^\infty_{2\pi}(\mathbb{R}^d, \mathbb{R})$, $r_1, \dots, r_d \in \mathbb{N}$, $m_1, \dots, m_d \in \mathbb{N}$, then we have*

$$\|f - P^{[1,\dots,d]}_{m_1,\dots,m_d}(f)\|_{C^0} \leq C_d \sum_{j=1}^{d} \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0}, \tag{3.3.5}$$

where C_d is a constant only depending on d.

Proof From [A], we will prove Lemma 3.4 by induction. The case $d = 1$ is covered by Jackson's approximation theorem. More precisely, for $f \in C^\infty_{2\pi}(\mathbb{R}, \mathbb{R})$, $m, r \in \mathbb{N}$, we have

$$\|f - P^{[1]}_m(f)\|_{C^0} \leq C_1 \frac{1}{m^r} \left\| \frac{\partial^r f}{\partial x^r} \right\|_{C^0}. \tag{3.3.6}$$

Let the assertion be true for some $d \in \mathbb{N}$. We verify it for $d+1$. Consider the functions $f(x_1, \cdot)$ with x_1 as a real parameter. Then by the assertion for d, we have

$$\|f(x_1, \cdot) - P^{[2,\dots,d+1]}_{m_2,\dots,m_{d+1}}(f)(x_1, \cdot)\|_{C^0} \leq C_d \sum_{j=2}^{d+1} \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0},$$

hence,

$$\|f - P^{[2,\dots,d+1]}_{m_2,\dots,m_{d+1}}(f)\|_{C^0} \leq C_d \sum_{j=2}^{d+1} \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0}. \tag{3.3.7}$$

Let $\hat{x}_j \in \mathbb{R}^d$ denote the vector $x \in \mathbb{R}^{d+1}$ without its j-th entry. For the functions $f(\cdot, \hat{x}_1)$, from (3.3.6), it follows that

$$\|f(\cdot, \hat{x}_1) - P^{[1]}_{m_1}(f)(\cdot, \hat{x}_1)\|_{C^0} \leq C_1 \frac{1}{{m_1}^{r_1}} \left\| \frac{\partial^{r_1} f}{\partial {x_1}^{r_1}} \right\|_{C^0},$$

hence,

$$\|f - P^{[1]}_{m_1}(f)\|_{C^0} \leq C_1 \frac{1}{{m_1}^{r_1}} \left\| \frac{\partial^{r_1} f}{\partial {x_1}^{r_1}} \right\|_{C^0}. \tag{3.3.8}$$

By (3.3.2), (3.3.3)),(3.3.4) and (3.3.7), we have

$$\begin{aligned}\left\| P^{[1]}_{m_1}(f) - P^{[1,\dots,d+1]}_{m_1,\dots,m_{d+1}}(f) \right\|_{C^0} &= \left\| P^{[1]}_{m_1}(f) - P^{[1]}_{m_1}\left(P^{[2,\dots,d+1]}_{m_2,\dots,m_{d+1}}(f)\right) \right\|_{C^0}, \\ &= \left\| P^{[1]}_{m_1}\left(f - P^{[1]}_{m_1} P^{[2,\dots,d+1]}_{m_2,\dots,m_{d+1}}(f)\right) \right\|_{C^0}, \\ &\leq 3C_d \sum_{j=2}^{d+1} \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0},\end{aligned}$$

which together with (3.3.8) implies that

$$\begin{aligned}\left\| f - P^{[1,\dots,d+1]}_{m_1,\dots,m_{d+1}}(f) \right\|_{C^0} &\leq \left\| f - P^{[1]}_{m_1}(f) \right\|_{C^0} + \left\| P^{[1]}_{m_1}(f) - P^{[1]}_{m_1}\left(P^{[2,\dots,d+1]}_{m_2,\dots,m_{d+1}}(f)\right) \right\|_{C^0}, \\ &\leq C_1 \frac{1}{{m_1}^{r_1}} \left\| \frac{\partial^{r_1} f}{\partial {x_1}^{r_1}} \right\|_{C^0} + 3C_d \sum_{j=2}^{d+1} \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0}, \\ &\leq C_{d+1} \sum_{j=1}^{d+1} \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0}.\end{aligned}$$

This finishes the proof. □

Obviously, there exist $m_{\bar{j}}, r_{\bar{j}}$ such that

$$\frac{1}{{m_{\bar{j}}}^{r_{\bar{j}}}} \left\| \frac{\partial^{r_{\bar{j}}} f}{\partial {x_{\bar{j}}}^{r_{\bar{j}}}} \right\|_{C^0} = \max_{1\leq j \leq d} \left\{ \frac{1}{{m_j}^{r_j}} \left\| \frac{\partial^{r_j} f}{\partial {x_j}^{r_j}} \right\|_{C^0} \right\}.$$

Hence, we have

$$\begin{aligned}\|f - P^{[1,\dots,d]}_{m_1,\dots,m_d}(f)\|_{C^0} &\leq dC_d \frac{1}{{m_{\bar{j}}}^{r_{\bar{j}}}} \left\| \frac{\partial^{r_{\bar{j}}} f}{\partial {x_{\bar{j}}}^{r_{\bar{j}}}} \right\|_{C^0}, \\ &\leq C'_d \frac{1}{{m_{\bar{j}}}^{r_{\bar{j}}}} \|f\|_{C^{r_{\bar{j}}}}.\end{aligned}$$

For the simplicity of notations, we denote

$$p_N(x) = P^{[1,\dots,d]}_{m_1,\dots,m_d}(f)(x),$$

where $x = (x_1, \dots, x_d)$ and $N = 2m_{\bar{j}} - 1$. Moreover, we denote $k := r_{\bar{j}}$, then

$$\|f(x) - p_N(x)\|_{C^0} \leq A_{dk} N^{-k} \|f(x)\|_{C^k}, \tag{3.3.9}$$

where A_{dk} is a constant depending on d and k.

3.4. C^ω case

Similar to Herman's construction, we consider C^∞ function $\tilde{T}_n(x)$ as follow:

$$\tilde{T}_n(x) = \begin{cases} \tilde{T}^+_n(x), & x \in [0,\pi]^d, \\ \tilde{T}^-_n(x), & x \in [-\pi, 0]^d, \\ 0, & \text{others.} \end{cases}$$

$\tilde{T}^+_n(x)$ satisfies:

$$\begin{cases} \operatorname{supp} \tilde{T}^+_n(x) \subset [0,\pi]^d, \\ \max \tilde{T}^+_n(x) \sim 1, \end{cases}$$

$\tilde{T}^-_n(x)$ satisfies:

$$\begin{cases} \operatorname{supp} \tilde{T}^-_n(x) = B_{R_n}(x_0), \\ \max \tilde{T}^-_n(x) = n, \\ R_n \sim \left(\frac{1}{n}\right)^{\frac{1}{d}}, \\ x_0 = \left(-\frac{\pi}{2}, \cdots, -\frac{\pi}{2}\right). \end{cases}$$

By Lemma 3.4, there exists a trigonometric polynomial $p_N(x_1, \cdots, x_d)$ in x_l $(1 \leq l \leq d)$ of degree at most N such that

$$\left\|\tilde{T}_n(x_1, \cdots, x_d) - p_N(x_1, \cdots, x_d)\right\|_{C^0} \leq A_{dk} N^{-k} \left\|\tilde{T}_n(x_1, \cdots, x_d)\right\|_{C^k}.$$

By the construction of $\tilde{T}_n$, we have

$$||\tilde{T}_n(x_1, \cdots, x_d)||_{C^k} \sim n^{\frac{k}{d}+1}. \tag{3.4.1}$$

Then, choosing N large enough such that

$$A_{dk} N^{-k} ||\tilde{T}_n(x_1, \cdots, x_d)||_{C^k} < \sigma \ll 1, \tag{3.4.2}$$

where σ is a small enough positive constant. Hence, we have

$$\begin{cases} \max p_N(x) \sim n, & \text{attained on } B_{R_n}(x_0), \\ \max p_N(x) \sim 1, & \text{on } [-\pi, \pi]^d \backslash \, B_{R_n}(x_0). \end{cases} \tag{3.4.3}$$

By (3.4.1) and (3.4.2), we have

$$N > \left(\frac{A_{dk}}{\sigma}\right)^{\frac{1}{k}} n^{\frac{1}{d}+\frac{1}{k}} \geq C n^{\frac{1}{d}+\frac{1}{k}}, \tag{3.4.4}$$

where C is a constant independent of n. We consider the normalized trigonometric polynomial

$$\tilde{p}_N(x) = \frac{1}{n^{1-\varepsilon}} \frac{p_N(x)}{\max |p_N(x)|}, \tag{3.4.5}$$

where $x = (x_1, \ldots, x_d)$. It is easy to see that

$$\begin{cases} \max \tilde{p}_N(x) \sim \frac{1}{n^{2-\varepsilon}}, \\ \min \tilde{p}_N(x) \sim -\frac{1}{n^{1-\varepsilon}}. \end{cases}$$

Hence, (3.2.3) is verified.

Next, we estimate $||\tilde{p}_N(x)||_{C^r}$. By a simple calculation, we have

$$||p_N(x)||_{C^r} \leq C n N^{r+1}. \tag{3.4.6}$$

Then,

$$\begin{aligned} ||\tilde{p}_N(x)||_{C^r} &\leq \frac{1}{n^{1-\varepsilon}} \frac{1}{\max |p_N(x)|} ||p_N(x)||_{C^r}, \\ &\leq \frac{1}{n^{2-\varepsilon}} \cdot C n N^{r+1}, \\ &= C \frac{1}{n^{1-\varepsilon}} N^{r+1}. \end{aligned}$$

To achieve $n^{-(1-\varepsilon)} N^{r+1} \to 0$ as $n \to \infty$, it suffices to make

$$\frac{1}{n^{1-\varepsilon}} N^{r+1} \leq \frac{1}{n^{\varepsilon}}.$$

Hence, we have

$$r \leq \log_N n^{1-2\varepsilon} - 1.$$

From (3.4.4), we take

$$N \sim n^{\frac{1}{d}+\frac{1}{k}},$$

then, we have

$$\max_N r \leq \left(\frac{1}{d}+\frac{1}{k}\right)^{-1}(1-2\varepsilon)-1. \tag{3.4.7}$$

Since k can be made large enough, we take $k=\frac{d}{2\varepsilon}$. Let

$$\delta = \frac{4\varepsilon}{1+2\varepsilon}d,$$

then, we have

$$\max_N r \leq d-1-\delta. \tag{3.4.8}$$

It follows that $\tilde{p}_N(x) \to 0$ as $n \to \infty$ in the $C^{d-1-\delta}$ for any $\delta > 0$.

Let $\tilde{\Psi}_n$ be the unique function in $C^{\omega}(\mathbb{T}^d, \mathbb{R})$ such that

$$\int_{\mathbb{T}^d} \tilde{\Psi}_n(x)dx = 0 \quad \text{and} \quad \frac{1}{d}\Delta\tilde{\Psi}_n(x) = \tilde{p}_N(x).$$

By Schauder estimates one knows that for any $\delta > 0$, $\tilde{\Psi}_n(x) \to 0$ as $n \to \infty$ in the $C^{d+1-\delta}$ topology. From the construction of $\tilde{p}_N(x)$, it is easy to see that (3.2.3) is verified. Hence, we have that the symplectic twist maps generated by the generating function $\tilde{h}_n(x,x') = \frac{1}{2}(x-x')^2 + \tilde{\Psi}_n(x')$ do not admit any Lagrangian tori for n large enough.

So far, we prove the following theorem

Theorem 3.5 *All Lagrangian tori of an integrable symplectic twist map with $d \geq 2$ degrees of freedom can be destructed by C^{ω} perturbations of the generating function and the perturbations are arbitrarily small in the $C^{d+1-\delta}$ topology for a small given constant $\delta > 0$.*

Based on the correspondence between symplectic twist maps and Hamiltonian systems, together with the toy model corresponding to the case with $d = 1$, we have the following corollary.

Corollary 3.6 *All Lagrangian tori of an integrable Hamiltonian system with $d \geq 2$ degrees of freedom can be destructed by C^{ω} perturbations which are arbitrarily small in the $C^{d-\delta}$ topology for a small given constant $\delta > 0$.*

Acknowledgement We would like to thank Prof. C.-Q. Cheng for many helpful discussions. This work is under the support of Research and Innovation Project for College Graduates of Jiangsu Province (CXZZ12_0030), the NNSF of China (Grant 11171146) and PAPD of Jiangsu Province of China.

References

[A] J.Albrecht. *On the existence of invariant tori in nearly-integrable Hamiltonian systems with finitely differentiable perturbations*. Regular and Chaotic Dynamics. **12** (2007), 281-320.

[Ba] V.Bangert. *Mather sets for twist maps and geodesics on tori*. Dynamics Reported **1** (1988), 1-45.

[Be] U.Bessi. *An analytic counterexample to KAM theorem*. Ergod. Th. & Dynam. Sys. **20** (2000), 317-333.

[BP] M.Bialy and L.Polterovich. *Hamiltonian systems, Lagrangian tori and Birkhoff's theorem*. Math. Ann. **292** (1992), 619-627.

[Ch] C.-Q.Cheng. *Non-existence of KAM torus*. Acta Mathmatica Sinica. **27** (2011), 397-404.

[Ci] B.V.Chirikov. *A universal instability of many-dimensional oscillator systems*. Phys. Reports. **52** (1979), 264-379.

[CW] C.-Q.Cheng and L.Wang. *Destruction of Lagrangian torus in positive definite Hamiltonian systems*. Preprint (2012).

[F] G.Forni. *Analytic destruction of invariant circles*. Ergod. Th. & Dynam. Sys. **14** (1994), 267-298.

[Go] C.Golé. *Optical Hamiltonians and symplectic twist maps*. Physica D: Nonlinear Phenomena. **71** (1994), 185-195.

[Gr] J.M.Greene *A method for determining a stochastic transition*. J.Math. Phys. **20** (1979), 1183-1201.

[H1] M.R.Herman. *Sur la conjugation différentiable des difféomorphismes du cercle à des rotations*. Publ. Math. IHES **49** (1979), 5-233.

[H2] M.R.Herman. *Sur les courbes invariantes par les difféomorphismes de l'anneau*. Astérisque **103-104** (1983), 1-221.

[H3] M.R.Herman. *Sur les courbes invariantes par les difféomorphismes de l'anneau*. Astérisque **144** (1986), 1-243.

[H4] M.R.Herman. *Inégalités "a priori" pour des tores lagrangiens invariants par des difféomrphismes symplectiques*. Inst. Hautes Études Sci. Publ. Math. **70** (1990), 47-101.

[H5] M.R.Herman. *Non existence of Lagrangian graphs*. unpublished preprint (1990).

[Ma1] J.N.Mather. *Existence of quasi periodic orbits for twist homeomorphisms of the annulus*. Topology **21** (1982), 457-467.

[Ma2] J.N.Mather. *Non-existence of invariant circles*. Ergod. Th. & Dynam. Sys. **4** (1984), 301-309.

[Ma3] J.N.Mather. *A criterion for the non-existence of invariant circle*. Publ. Math. IHES **63** (1986), 301-309.

[Ma4] J.N.Mather. *Modulus of continuity for Peierls's barrier*. Periodic Solutions of Hamiltonian Systems and Related Topics. ed. P.H.Rabinowitz *et al.* NATO ASI Series C **209**. Reidel: Dordrecht, (1987), 177-202.

[Ma5] J.N.Mather. *Destruction of invariant circles*. Ergod. Th. & Dynam. Sys. **8** (1988), 199-214.

[Ma6] J.N.Mather. *Action minimizing invariant measures for postive definite Lagrangian systems*. Math. Z. **207** (1991), 169-207.

[Mo1] J.Moser. *On invariant curves of area-preserving mappings of an annulus*. Nachr. Akad. Wiss. Göttingen, II. Math-Phys. KL. **1** (1962), 1-20.

[Mo2] J.Moser. *A rapidly convergent iteration method and non-linear partail differential equations I, II.* Annali della Scuola Normale Superiore di Pisa. **20** (1966), 265-315.

[Mo3] J.Moser. *On the construction of almost periodic solutions for ordinary differential equations.* Proceedings of the International Conference on Functional Analysis and Related Topics, Tokyo. (1969), 60-67.

[Mo4] J.Moser. *Monotone twist mappings and the calculus of variations.* Ergod. Th. & Dynam. Sys. **6** (1986), 401-413.

[MP] R.S.MacKay and I.C.Percival *Converse KAM: theory and practice.* Commun. Math. Phys. **98** (1985), 469-512.

[P] J.Pöschel. *Integrability of Hamiltonian systems on Cantor sets.* Comm. Pure Appl. Math. **35** (1982), 653-696.

[R1] H.Rüssman. *On optimal estimates for the solutions of linear partial differential equations of first order with constant coefficients on the torus.* Lecture Notes in Physics. **38** (1975), 598-624.

[R2] H.Rüssman. *On the existence of invariant curves of twist mappings of an annulus.* Lecture Notes in Mathematics. **1007** (1983), 677-718.

[W1] L.Wang. *Variational destruction of invariant circles.* Discrete and Continuous Dynamical Systems-A. **32** (2012), 4429-4443.

[W2] L.Wang. *Analytic destruction of invariant circles for exact area-preserving twist maps.* Preprint (2012).

[W3] L.Wang. *Analytic destruction of Lagrangian torus.* Preprint (2012).

[W4] L.Wang. *Total destruction of Lagrangian tori.* Preprint (2012).

[Z] A.Zygmund. *Trigonometric Series.* Third Edition Volumes I & II combined, with a foreword by Robert Fefferman. Cambridge University Press, Cambridge, 2002.

Department of Mathematics, Nanjing University, Nanjing 210093, China.

E-mail address: `linwang.math@gmail.com`